

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
29 March 2001 (29.03.2001)

PCT

(10) International Publication Number  
WO 01/21768 A1

(51) International Patent Classification<sup>7</sup>: C12N 5/04,  
15/09, 15/63, 15/64, 15/65, 15/67, 15/82, 15/87, 15/90,  
A01H 1/00, 5/00

Street, Apt. A, Highland Park, NJ 08904 (US). LUTZ,  
Kerry [US/US]; 8 Liberty Place, Jackson, NJ 08527 (US).

(21) International Application Number: PCT/US00/25930

(74) Agents: RIGAUT, Kathleen, D. et al.; Dann, Dorfman,  
Herrell and Skillman, Suite 720, 1601 Market Street,  
Philadelphia, PA 19120 (US).

(22) International Filing Date:  
21 September 2000 (21.09.2000)

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,  
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,  
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,  
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,  
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,  
TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/155,007 21 September 1999 (21.09.1999) US  
60/211,139 13 June 2000 (13.06.2000) US

(84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian  
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European  
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,  
IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG,  
CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): RUT-  
GERS, THE STATE UNIVERSITY OF NEW JERSEY  
[US/US]; Old Queens, Somerset Street, New Brunswick,  
NJ 08903 (US).

Published:  
— With international search report.

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): MALIGA, Pal  
[HU/US]; 72 Yorktown Road, East Brunswick, NJ 08816  
(US). CORNEILLE, Sylvie [FR/US]; 305 Magnolia

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.



WO 01/21768 A1

(54) Title: SITE-SPECIFIC RECOMBINATION SYSTEM TO MANIPULATE THE PLASTID GENOME OF HIGHER PLANTS

(57) Abstract: A site specific recombination system and methods of use thereof are disclosed for manipulating the genome of higher plants.

**Site-specific recombination system to manipulate the  
plastid genome of higher plants**

5

10           This application claims priority under 35 U.S.C.  
§119(e) to US Provisional Applications 60/155,007 and  
60/211,139 filed September 21, 1999 and June 13, 2000  
respectively, the entire disclosure of each of the  
above-identified applications is incorporated by  
reference herein.

15

**FIELD OF THE INVENTION**

20           This invention relates to the fields of transgenic  
plants and molecular biology. More specifically, DNA  
constructs and methods of use thereof are provided which  
facilitate the excision of target DNA sequences from  
transplastomic plants.

**BACKGROUND OF THE INVENTION**

25           Several publications are referenced in this  
application by author name and year of publication in  
parentheses in order to more fully describe the state of  
the art to which this invention pertains. Full  
citations for these reference can be found at the end of  
30           the specification. The disclosure of each of these  
publications is incorporated by reference herein.

35           The plastid genetic system of higher plants is  
highly polyploid. For example, in a tobacco leaf there  
are as many as 100 chloroplasts, each carrying ~100  
identical genome copies, a total of 10,000 copies in a  
leaf cell. High-level protein expression, lack of  
pollen transmission and the feasibility to engineer  
polycistronic expression units make the plastid genome

an attractive alternative to nuclear engineering.

Plastid transformation vectors often contain a selective marker, most commonly a spectinomycin resistance (*aadA*) gene, flanked by plastid DNA sequences targeting

5 insertion of the marker gene by homologous recombination into the plastid genome. Genes of commercial value but lacking a selectable phenotype are physically linked to the selective marker and the two genes are integrated together as a block of heterologous sequences. Plastid  
10 transformation is accomplished by biolistic DNA delivery or polyethylene glycol induced uptake of the transforming DNA followed by selection for the antibiotic resistance marker to ensure preferential propagation of plastids with transformed genome copies.  
15 As the result, all the 10,000 wild-type plastid genome copies in a cell are replaced with transgenic copies during a gradual process (Maliga, 1993).

Incorporation of a selectable marker gene is essential to ensure preferential maintenance of the  
20 transformed plastid genome copies. However, once transformation is accomplished, maintenance of the marker gene is undesirable. One problem may be the metabolic burden imposed by the expression of the selectable marker gene. For example FLARE-S, the product  
25 of the marker gene with good prospects to transform cereal chloroplasts, accumulates up to 18% of the total soluble cellular protein (Khan and Maliga 1999). The second problem is the relatively high potential for horizontal transfer of plastid marker genes to microbes  
30 (Tepfer 1989; Dröge et al. 1998; Sylvanen 1999), as commonly used plastid marker gene constructs are efficiently expressed in *E. coli* (Carrer et al. 1993; Svab and Maliga 1993). Therefore, having plastid marker genes in commercial products is undesirable.

35

**SUMMARY OF THE INVENTION**

In accordance with the present invention, methods and systems are provided which facilitate the manipulation of the plastid genomes of higher plants. The methods and systems of the invention may be employed to remove heterologous sequences from the plastid genome, such as selectable marker genes following successful isolation of transformed progeny. Alternatively, they may be designed to remove endogenous genes involved in plant cell metabolism, growth, development and fertility.

In one embodiment of the invention, a site specific recombination method for removal of predetermined nucleic acid sequences from the plastid genome is provided. The method comprises providing a first nucleic acid construct, the construct comprising a promoter being operably linked to a nucleic acid encoding an optional plastid targeting transit sequence which is in turn operably linked to a nucleic acid encoding a protein having excision activity, the construct further comprising a first selectable marker encoding nucleic acid having plant specific 5' and 3' regulatory nucleic acid sequences. The method also entails the use of a second DNA construct, the second construct comprising an second selectable marker encoding nucleic acid and excision sites. The second construct optionally contains a gene of interest and further comprises flanking plastid targeting nucleic acid sequences which facilitate homologous recombination into said plastid genome. The second DNA construct is introduced into plant cell and the cells are cultured in the presence of a selection agent, thereby selecting for those plant cells expressing the proteins encoded by said second DNA construct. The first DNA construct is

then introduced into cells having the second construct in the presence of a selection agent and those plant cells expressing proteins encoded by said first construct are selected. If present, the excising activity acts on the excision sites, thereby excising said predetermined target sequence. Plants may then be regenerated from plant cells obtained by the foregoing method.

Proteins having excision activity suitable for the practice of the invention include, without limitation, CRE, flippase, resolvase, FLP, SSV1-encoded integrase, and transposase. Sequences corresponding to excision sites suitable for the practice of the invention, include, for example, LOX sequences, and *frt* sequences.

A variety of selection of agents may be selected. These include without limitation, kanamycin, gentamycin, spectinomycin, streptomycin and hygromycin, phosphinotricin, basta, glyphosate and bromoxynil.

In an alternative embodiment, a site specific recombination method for removal of predetermined nucleic acid sequences from the plastid genome is provided. The method comprising providing a first nucleic acid construct, said construct comprising a regulated promoter being operably linked to a nucleic acid encoding an optional plastid targeting transit sequence which is operably linked to a nucleic acid encoding a protein having excision activity, said construct optionally further comprising a first selectable marker encoding nucleic acid having plant specific 5' and 3' regulatory nucleic acid sequences. A second DNA construct is also provided, said second construct comprising an second selectable marker encoding nucleic acid and excision sites, said second construct further comprising flanking plastid targeting nucleic acid sequences which facilitate homologous

recombination into said plastid genome at a predetermined target sequence such that excision sites flank said predetermined target sequence following homologous recombination and introducing said second DNA construct into a plant cell. The plant cell so generated is then cultured in the presence of a selection agent, thereby selecting for those plant cells expressing the proteins encoded by said second DNA construct. A plant is then regenerated from cells containing the second construct and the first DNA construct is introduced into these cells in the presence of a selection agent and those plant cells expressing proteins encoded by said first construct are selected. The excising activity then acts on the excision sites, thereby excising said predetermined target sequence.

Regulatable promoters suitable for this embodiment of the invention include, without limitation, inducible promoters, tissue specific promoters, developmentally regulated promoters and chemically inducible promoters.

Candidate predetermined target sequences, may include for example genes associated with male sterility, *clpP*, ribosomal proteins, ribosomal operon sequences.

## BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is a schematic diagram depicting CRE-mediated excision and integration of DNA segments.

Figure 2 is a map of a plastid transformation vector pSAC48, with *codA* bracketed by direct *loxP* sites. Positions of plastid genes *rrn16*, *trnV*, *rps12/7* (Shinozaki et al. 1986), the *aadA* and *codA* transgenes and relevant restriction sites are marked.

Figure 3 is a map of an *Agrobacterium* binary vector

pPZP212 with a plastid-targeted *Ssu-tp-cre* gene. Marked are: *Agrobacterium* Left and Right Border fragments; the kanamycin resistance (*neo*) gene; P2' promoter; SSU transit peptide (*ssu-tp*); *cre* coding region; recognition sequences for restriction enzymes *Bam*HI, *Eco*RI, *Hind*III, *Nco*I, *Nhe*I and *Xba*I.

Figure 4 shows maps of the plastid genome >*codA*> deletion derivatives. Shown are the plastid targeting region of vector pSAC48; the map of same region of the wild-type plastid genome (Nt-wt); the map of the plastid genome with CRE-mediated deletion of *codA* via the *lox* sites; and the map of the plastid genome with deletion via *Prn* sequences lacking *trnV*, *aadA* and *codA*. Positions of plastid genes *rrn16*, *trnV* and *rps12/7* (Shinozaki et al. 1986), *aadA* and *codA* transgenes, primers (O1-O4) and relevant restriction sites (AI, *Apa*I; EV, *Eco*RV) are marked.

Figure 5 is a gel showing PCR amplification which confirms CRE-mediated deletion of *codA* from the plastid genome. Primers O1 and O2 (Fig. 3) amplified the 0.7-kb fragment of the deleted region. Same primers amplify the 2.0-kb *aadA-codA* fragment in tester lines Nt-pSAC48-21A and Nt-pSAC-16C (no transgenic Cre gene). No specific fragment was obtained in wild-type DNA sample and in Cre1-10 line. The lines obtained are listed in Table 1.

Figure 6 shows the results of DNA gel blot analysis wherein plastid genome structure was determined in the indicated plant samples. Total cellular DNA was isolated from the leaves of plants listed in Table 1 and digested with the *Apa*I and *Eco*RV restriction endonucleases. The probes were the wild-type *Apa*I-*Eco*RV plastid targeting

region and the *aadA* (*NcoI*-*XbaI* fragment) and *codA* (*NcoI*-*XbaI* fragment) coding regions. The hybridizing fragments are marked in Fig. 3.

5           Figure 7 are gels showing uniformity of plastid genome populations in the *Ssu-tp-cre* transformed plants. Total cellular DNA extracted from several leaves was probed with the *ApaI*-*EcoRV* targeting region probe. Numbers identify leaves from which DNA was extracted. 10           For example, seven different leaves were probed from the *Cre1-3* plant. For details, see Brief Description of Fig. 6.

          Figures 8A and 8B are gels of PCR analysis 15           confirming CRE-mediated deletion of *codA* in seedlings obtained by pollination with *Ssu-tp-cre* activator lines. 5-day old seedlings were tested from the cross *Nt-pSAC48-21A* as maternal parent and *Cre2-200* and *Cre2-300* activator lines as pollen parents. Amplification 20           products are also shown for controls *Nt-pSAC48-21A* selfed seedling (48 self), wild-type (wt), the parental plant (48P) and the *Cre1-3* plant. Fig. 8A: The *codA* region was amplified with the 01/02 primers: the size of *aadA-codA* fragment is 2.0 kb; the *codA* deletion fragment 25           is 0.7 kb (Fig. 4). Fig. 8B: Testing for *cre* sequences by PCR amplification with the *Cre1/Cre3* oligonucleotides.

          Figure 9 is a diagram of the plastid transformation 30           

*pSAC38* with the >neo< bracketed by inverted lox sites. Positions of plastid genes *rrn16*, *trnV* and *rps12/7* (Shinozaki et al., 1986), the *aadA* and *codA* transgenes and relevant restriction sites are marked.

35           Figure 10 shows a map of the plastid genome



containing the >neo< inversion construct. Shown are the plastid targeting region of vector pSAC38; the map of the same region of the wild-type plastid genome (Nt-wt); map of the plastid genome with CRE-mediated inversion of neo via the lox sites. Positions of the plastid genes *rrn16*, *trnV* and *rps12/7* (Shinozaki et al., 1986) *aadA* and *neo* transgenes, primers (01-04) and relevant restriction sites (BamHI) are marked.

Figure 11 shows the results of DNA gel blot analysis for the determination of plastid genome structure of CRE-activated >neo< plants by DNA gel blot analysis. Total cellular DNA was digested with the BamHI restriction endonuclease. The probes was the wild-type *ApaI-EcoRV* plastid targeting region. The hybridizing fragments are marked in Fig. 10.

Figure 12 shows an exemplary monocistronic inversion vector. The gene of interest (*goi*) coding region is flanked by inverted lox sites (triangles). CRE activates *goi* expression by inversion, so that the coding strand is transcribed. *rrn16*, *trnV* and *rps12/7* are plastid genes (Shinozaki et al. 1986).

Figure 13 shows an alternative dicistronic lox inversion vector. Note that the inverted lox sites flank the selective marker (*aadA*) and *goi*, and only one gene is expressed. *rrn16*, *trnV* and *rps12/7* are plastid genes (Shinozaki et al. 1986).

Figure 14 shows a basic tobacco plastid lox deletion vector. The vector provides is a suitable backbone for vector construction and targets insertions into the *trnV-rps12/7* intergenic region.

Figure 15 shows a tobacco plastid *lox* >*aadA*> deletion vector. *rrn16*, *trnV* and *rps12/7* are plastid genes (Shinozaki et al. 1986).

5

Figure 16 shows a tobacco constitutive >*aadA*>*goi* dicistronic deletion vector. *rrn16*, *trnV* and *rps12/7* are plastid genes and are described in (Shinozaki et al. 1986).

10

Figure 17 shows a tobacco constitutive *goi* >*aadA*> dicistronic deletion vector. Note that vectors shown in Fig. 16 and Fig. 17 differ in the relative order of marker gene and the gene of interest. *rrn16*, *trnV* and *rps12/7* are plastid genes (Shinozaki et al. 1986).

15

Figure 18 shows a tobacco constitutive *goi* >*aadA*> dicistronic deletion vector, in which expression of *aadA* is dependent on translational coupling. Note that in this construct only one leader sequence is utilized. *rrn16*, *trnV* and *rps12/7* are plastid genes (Shinozaki et al. 1986).

20

Figure 19 shows a tobacco inducible *lox* deletion vector. Expression of *goi* is dependent on *aadA* excision. *rrn16*, *trnV* and *rps12/7* are plastid genes (Shinozaki et al. 1986). Abbreviations: P, promoter; T, 3' untranslated region; L1 is 5' leader sequence.

25

Figure 20 shows a vector suitable for Cre-mediated deletion of *clpP* gene from the plastid genome. The region of engineered plastid genome shown is the sequence contained in the plastid transformation vector. The *clpP* Exons are dark boxes, the Introns are open

30

boxes. Map position of plastid genes *psbB*, *rps12* Exon I and *rpl20* is also shown.

#### DETAILED DESCRIPTION OF THE INVENTION

5       The following definitions are provided to aid in understanding the subject matter regarded as the invention.

10       Heteroplastomic refers to the presence of a mixed population of different plastid genomes within a single plastid or in a population of plastids contained in plant cells or tissues.

15       Homoplastomic refers to a pure population of plastid genomes, either within a plastid or within a population contained in plant cells and tissues.

20       Homoplastomic plastids, cells or tissues are genetically stable because they contain only one type of plastid genome. Hence, they remain homoplastomic even after the selection pressure has been removed, and selfed progeny are also homoplastomic. For purposes of the present invention, heteroplastomic populations of genomes that are functionally homoplastomic (i.e., contain only minor populations of wild-type DNA or transformed genomes with sequence variations) may be referred to herein as "functionally homoplastomic" or "substantially

25       homoplastomic." These types of cells or tissues can be readily purified to a homoplastomic state by continued selection.

      Plastome refers to the genome of a plastid.

30       Transplastome refers to a transformed plastid genome.

      Transformation of plastids refers to the stable integration of transforming DNA into the plastid genome that is transmitted to the seed progeny of plants containing the transformed plastids.

35       Selectable marker gene refers to a gene that upon

expression confers a phenotype by which successfully transformed plastids or cells or tissues carrying the transformed plastid can be identified.

Transforming DNA refers to homologous DNA, or  
5 heterologous DNA flanked by homologous DNA, which when introduced into plastids becomes part of the plastid genome by homologous recombination.

Operably linked refers to two different regions or two separate genes spliced together in a construct such  
10 that both regions will function to promote gene expression and/or protein translation.

"Nucleic acid" or a "nucleic acid molecule" as used herein refers to any DNA or RNA molecule, either single or double stranded and, if single stranded, the molecule  
15 of its complementary sequence in either linear or circular form. In discussing nucleic acid molecules, a sequence or structure of a particular nucleic acid molecule may be described herein according to the normal convention of providing the sequence in the 5' to 3'  
20 direction. With reference to nucleic acids of the invention, the term "isolated nucleic acid" is sometimes used. This term, when applied to DNA, refers to a DNA molecule that is separated from sequences with which it is immediately contiguous in the naturally occurring  
25 genome of the organism in which it originated. For example, an "isolated nucleic acid" may comprise a DNA molecule inserted into a vector, such as a plasmid or virus vector, or integrated into the genomic DNA of a prokaryotic or eukaryotic cell or host organism.

30 When applied to RNA, the term "isolated nucleic acid" refers primarily to an RNA molecule encoded by an isolated DNA molecule as defined above. Alternatively, the term may refer to an RNA molecule that has been sufficiently separated from other nucleic acids with  
35 which it would be associated in its natural state (i.e.,

in cells or tissues). An isolated nucleic acid (either DNA or RNA) may further represent a molecule produced directly by biological or synthetic means and separated from other components present during its production.

5           The terms "percent similarity", "percent identity" and "percent homology" when referring to a particular sequence are used as set forth in the University of Wisconsin GCG software program.

10           The term "functional" as used herein implies that the nucleic or amino acid sequence is functional for the recited assay or purpose.

15           The phrase "consisting essentially of" when referring to a particular nucleotide or amino acid means a sequence having the properties of a given SEQ ID No.:. For example, when used in reference to an amino acid sequence, the phrase includes the sequence per se and molecular modifications that would not affect the basic and novel characteristics of the sequence.

20           A "replicon" is any genetic element, for example, a plasmid, cosmid, bacmid, phage or virus, that is capable of replication largely under its own control. A replicon may be either RNA or DNA and may be single or double stranded.

25           A "vector" is a replicon, such as a plasmid, cosmid, bacmid, phage or virus, to which another genetic sequence or element (either DNA or RNA) may be attached so as to bring about the replication of the attached sequence or element.

30           An "expression operon" refers to a nucleic acid segment that may possess transcriptional and translational control sequences, such as promoters, enhancers, translational start signals (e.g., ATG or AUG codons), polyadenylation signals, terminators, and the like, and which facilitate the expression of a  
35           polypeptide coding sequence in a host cell or organism.

The term "oligonucleotide," as used herein refers to primers and probes of the present invention, and is defined as a nucleic acid molecule comprised of two or more ribo- or deoxyribonucleotides, preferably more than  
5 three. The exact size of the oligonucleotide will depend on various factors and on the particular application and use of the oligonucleotide.

The term "probe" as used herein refers to an oligonucleotide, polynucleotide or nucleic acid, either  
10 RNA or DNA, whether occurring naturally as in a purified restriction enzyme digest or produced synthetically, which is capable of annealing with or specifically hybridizing to a nucleic acid with sequences complementary to the probe. A probe may be either  
15 single-stranded or double-stranded. The exact length of the probe will depend upon many factors, including temperature, source of probe and use of the method. For example, for diagnostic applications, depending on the complexity of the target sequence, the oligonucleotide  
20 probe typically contains 15-25 or more nucleotides, although it may contain fewer nucleotides. The probes herein are selected to be "substantially" complementary to different strands of a particular target nucleic acid sequence. This means that the probes must be  
25 sufficiently complementary so as to be able to "specifically hybridize" or anneal with their respective target strands under a set of pre-determined conditions. Therefore, the probe sequence need not reflect the exact complementary sequence of the target. For example, a  
30 non-complementary nucleotide fragment may be attached to the 5' or 3' end of the probe, with the remainder of the probe sequence being complementary to the target strand. Alternatively, non-complementary bases or longer sequences can be interspersed into the probe, provided  
35 that the probe sequence has sufficient complementarity

with the sequence of the target nucleic acid to anneal therewith specifically.

The term "primer" as used herein refers to an oligonucleotide, either RNA or DNA, either  
5 single-stranded or double-stranded, either derived from a biological system, generated by restriction enzyme digestion, or produced synthetically which, when placed in the proper environment, is able to functionally act as an initiator of template-dependent nucleic acid  
10 synthesis. When presented with an appropriate nucleic acid template, suitable nucleoside triphosphate precursors of nucleic acids, a polymerase enzyme, suitable cofactors and conditions such as a suitable temperature and pH, the primer may be extended at its 3'  
15 terminus by the addition of nucleotides by the action of a polymerase or similar activity to yield an primer extension product. The primer may vary in length depending on the particular conditions and requirement of the application. For example, in diagnostic  
20 applications, the oligonucleotide primer is typically 15-25 or more nucleotides in length. The primer must be of sufficient complementarity to the desired template to prime the synthesis of the desired extension product, that is, to be able anneal with the desired template  
25 strand in a manner sufficient to provide the 3' hydroxyl moiety of the primer in appropriate juxtaposition for use in the initiation of synthesis by a polymerase or similar enzyme. It is not required that the primer sequence represent an exact complement of the desired  
30 template. For example, a non-complementary nucleotide sequence may be attached to the 5' end of an otherwise complementary primer. Alternatively, non-complementary bases may be interspersed within the oligonucleotide primer sequence, provided that the primer sequence has  
35 sufficient complementarity with the sequence of the

desired template strand to functionally provide a template-primer complex for the synthesis of the extension product. Amino acid residues described herein are preferred to be in the "L" isomeric form. However, residues in the "D" isomeric form may be substituted for any L-amino acid residue, provided the desired properties of the polypeptide are retained.

All amino-acid residue sequences represented herein conform to the conventional left-to-right amino-terminus to carboxy-terminus orientation.

The term "tag," "tag sequence" or "protein tag" refers to a chemical moiety, either a nucleotide, oligonucleotide, polynucleotide or an amino acid, peptide or protein or other chemical, that when added to another sequence, provides additional utility or confers useful properties, particularly in the detection or isolation, to that sequence. Thus, for example, a homopolymer nucleic acid sequence or a nucleic acid sequence complementary to a capture oligonucleotide may be added to a primer or probe sequence to facilitate the subsequent isolation of an extension product or hybridized product. In the case of protein tags, histidine residues (e.g., 4 to 8 consecutive histidine residues) may be added to either the amino- or carboxy-terminus of a protein to facilitate protein isolation by chelating metal chromatography. Alternatively, amino acid sequences, peptides, proteins or fusion partners representing epitopes or binding determinants reactive with specific antibody molecules or other molecules (e.g., flag epitope, c-myc epitope, transmembrane epitope of the influenza A virus hemagglutinin protein, protein A, cellulose binding domain, calmodulin binding protein, maltose binding protein, chitin binding domain, glutathione S-transferase, and the like) may be added to proteins to



facilitate protein isolation by procedures such as affinity or immunoaffinity chromatography. Chemical tag moieties include such molecules as biotin, which may be added to either nucleic acids or proteins and facilitates isolation or detection by interaction with avidin reagents, and the like. Numerous other tag moieties are known to, and can be envisioned by, the trained artisan, and are contemplated to be within the scope of this definition.

As used herein, the terms "reporter," "reporter system", "reporter gene," or "reporter gene product" shall mean an operative genetic system in which a nucleic acid comprises a gene that encodes a product that when expressed produces a reporter signal that is a readily measurable, e.g., by biological assay, immunoassay, radioimmunoassay, or by colorimetric, fluorogenic, chemiluminescent or other methods. The nucleic acid may be either RNA or DNA, linear or circular, single or double stranded, antisense or sense polarity, and is operatively linked to the necessary control elements for the expression of the reporter gene product. The required control elements will vary according to the nature of the reporter system and whether the reporter gene is in the form of DNA or RNA, but may include, but not be limited to, such elements as promoters, enhancers, translational control sequences, poly A addition signals, transcriptional termination signals and the like.

The terms "transform", "transfect", "transduce", shall refer to any method or means by which a nucleic acid is introduced into a cell or host organism and may be used interchangeably to convey the same meaning. Such methods include, but are not limited to, transfection, electroporation, microinjection, PEG-fusion, biolistic bombardment and the like.

A "clone" or "clonal cell population" is a population of cells derived from a single cell or common ancestor by mitosis.

5 A "cell line" is a clone of a primary cell or cell population that is capable of stable growth *in vitro* for many generations.

#### CRE-MEDIATED SITE SPECIFIC RECOMBINATION

10 The plastid genome of higher plants is present in 100-10,000 copies per cell. Incorporation of a selectable marker gene is essential to ensure preferential maintenance of the transformed plastid genome copies carrying useful genes with no selectable phenotype. However, once transformation is accomplished, 15 maintenance of the marker gene is undesirable. In accordance with the present invention, a bacteriophage P1CRE-loxP site-specific recombination system is provided which is suitable for efficient elimination of marker genes from the plastid genome. The system 20 exemplified herein has two components: a plastid tester strain carrying a cytosine deaminase (*codA*) transgene flanked by lox sites conferring sensitivity to 5-fluorocytosine and a nuclear CRE line carrying a nuclear-encoded, plastid-targeted CRE. Both the plastid 25 tester (no CRE activity) and the nuclear CRE line (no lox sequence) were genetically stable. However, *codA* was eliminated at a very fast rate when the plastid-targeted CRE was introduced into the plastid tester strain by transformation or crossing. The gene for the nuclear- 30 encoded CRE was subsequently separated from the transformed plastids by segregation in the seed progeny. Excision of *codA* by CRE was often accompanied by deletion of a plastid genome segment flanked by short directly repeated sequences. Removal of the antibiotic 35 resistance marker from the transplastomic plants

eliminates the metabolic burden imposed by the expression of the selectable marker gene and should also improve public acceptance of the transgenic crops.

Additional applications of the CRE-lox site-specific recombination system are activation of plastid gene expression by deletion or inversion of plastid genome sequences and induction of controlled cell death by deleting vital genes in the male reproductive tissue.

Although the use the CRE recombinase is exemplified herein, other prokaryotic and eukaryotic site-specific recombinases would be equally suitable for the elimination of the marker genes.

Recently, several prokaryotic and lower eukaryotic site-specific recombination systems have been shown to operate successfully in higher eukaryotes. In plant and animal cells functional site-specific recombination systems from bacteriophages P1 (Cre-lox) Mu (Gin-gix), and from the inversion plasmids of *Saccharomyces cerevisiae* (FLP-*frt*) (Morris et al. 1991; O'Gorman et al. 1991; Lichtenstein and Barrena 1993; Lyznik et al. 1993; Lyznik et al. 1995; Lyznik et al. 1996) and *Zygosaccharomyces rouxii* (R-RS). In each of these systems, no additional factor aside from the recombinase and target sequences is required for recombination.

Reviewed in van Haaren and Ow, 1993. The CRE-loxP site-specific recombination system of bacteriophage P1 has been studied extensively *in vitro* and in *E. coli* (Craig 1988; Adams et al. 1992). Expression of the CRE protein (38.5 kDa) is sufficient to cause recombination between 34 bp loxP sites that consist of 13 bp inverted repeats separated by 8 bp asymmetric spacer sequence. If there are two loxP sites within a DNA segment, the result of the recombination reaction depends on the relative position of the recombination sites. If the recombination sites form a direct repeat, that if they

are in the same orientation, recombination results in deletion of the intervening DNA. If the recombination sites are in an inverted orientation, CRE-mediated recombination results in an inversion of the intervening DNA. The products of these reactions are shown in Fig. 1. The CRE site-specific recombination system has been employed for the elimination of nuclear genes in a number of eukaryotic systems, including higher plants (Dale and Ow 1991; Russell et al. 1992; Srivastava et al. 1999).

Before the present invention, the efficiency of CRE-mediated elimination of targeted plastid genes was unknown. To explore this system for this purpose, CRE-mediated elimination of the *codA* gene encoding cytosine deaminase (CD; EC 3.5.4.1) was assessed. Cytosine deaminase converts 5-fluorocytosine (5FC) into 5-fluorouracil (5FU), the precursor of 5-fluoro-dUMP. 5FC is lethal for CD-expressing cells due to irreversible inhibition of thymidylate synthase by 5-fluoro-dUMP (Beck et al. 1972). Cytosine deaminase is absent in plants. Expression of the bacterial *codA* in plastids renders cells sensitive to 5FC, while cells deficient in transgene expression are resistant (Serino and Maliga 1997). Thus, 5FC resistance could be used for positive identification of cells with CRE-induced *codA* deletion, even if such deletion events were relatively rare. The test system of the present invention incorporates a *codA* gene in the tobacco plastid genome between two directly oriented *lox* sites (>*codA*>). The transplastome was stable in the absence of CRE activity. However, highly efficient elimination of >*codA*> was triggered by introduction of a nuclear-encoded plastid-targeted CRE.

#### EXAMPLE 1

**CRE-MEDIATED DELETION OF THE SELECTABLE PLASTID MARKER**

Cre-mediated deletion of the selective plastid marker in the plastids of tobacco somatic cell is described in Example I. The selectable marker flanked by the lox sites is exemplified here by *codA*. However, it could be any other selectable and non-selectable marker gene, or any DNA sequence independent of information content flanked by lox sites in the plastid genome. Components of the test system are tobacco plants carrying a *codA* coding region flanked by lox sites (>*codA*>). A second component of the test system is a nuclear gene encoding a plastid targeted CRE-site specific recombinase. Deletion of a plastid encoded >*codA*> is achieved by introducing nuclear Cre into the nucleus of somatic (leaf) tobacco cells by Agrobacterium-mediated transformation. Alternatively, the nuclear encoded Cre gene may be introduced by fertilization with pollen of an appropriate activator-of-deletion strain. The nuclear Cre gene is subsequently removed by segregation in the seed progeny.

**MATERIALS AND METHODS FOR THE PRACTICE OF EXAMPLE 1**

The following materials and methods are provided to facilitate the practice of Example 1.

**Plastid *codA* with direct lox sites.**

The *codA* gene is contained in a *SacI-HindHIII* fragment. The gene map is shown in Fig. 2. *PrnloxD* (Seq. ID No. 4) is a plastid rRNA operon (*rrn16*) promoter derivative. It is contained in a *SacI-EcoRI* fragment obtained by PCR using oligonucleotides 5'-  
GGGGAGCTCGCTCCCCCGCCGTCGTTCAATG-3' and 5'-  
GGGAATTCATAACTTCGTATAGCATACATTATACGAAGTTAT

GCTCCCAGAAATATAGCCA-3' as primers and plasmid pZS176 (progenitor of plasmid pZS197; Svab and Maliga 1993) as a template. The promoter fragment *PrrnloxD* contains a *lox* site at the 3' end adjacent to the *EcoRI* site. The *EcoRI-NcoI* fragment contains the ribosome binding site from plasmid pZS176. The fragment was obtained by annealing the complementary oligonucleotides 5'-AATTCGAAGCGCTTGGATACAGTTGTAGGGAGGGATC-3' and 5'-CATGGATCCCTCCCTACAACGTATCCAAGCGCTTCG-3'. The *codA* coding region is contained in an *NcoI-XbaI* fragment (Serino and Maliga 1997). The *TrbcLloxD* (Seq. ID No. 5) is the *rbcL* 3'-untranslated region contained in an *XbaI-HindIII* fragment obtained by PCR using oligonucleotides 5'-GGTCTAGATAACTTCGTATAATGTATGCTATA CGAAGTTATAGACATTAGCAGATAAATT-3' and 5'-GGGGGTACCAAGCTTGCTAGATTTTGTATTTCAAATCTTG-3' and plasmid pMSK48 (Khan and Maliga 1999) as template. *TrbcLloxD* contains a *lox* site adjacent to the *XbaI* site in direct orientation relative to the *lox* site in the *codA* 5'UTR. The chimeric *PrrnloxD:codA:TrbcLloxD* gene was introduced into the tobacco plastid transformation vector pPRV111B (Zoubenko et al. 1994) as a *SacI-HindIII* fragment to obtain plasmid pSAC48.

**Plastid-targeted nuclear *cre* linked to a nuclear kanamycin resistance gene.** Two plastid targeted nuclear *cre* genes were tested. The *cre* gene in *Agrobacterium* binary vector pK027 and pK028 encode the CRE recombinase at its N terminus translationally fused with the pea Rubisco small subunit (SSU) chloroplast transit peptide (Timko et al. 1985) and twenty two and five amino acids of the mature Rubisco small subunit, respectively. Both *cre* genes are contained in an *EcoRI-HindIII* fragment. The schematic map of the genes is shown in Fig. 3. The

P2' *Agrobacterium* promoter (Velten et al. 1984) (Sequence ID. No.9) is contained in an *EcoRI*-*NcoI* fragment. The P2' promoter fragment was obtained by PCR using oligonucleotides 5'-

5     ccgaattcCATTTTCACGTGTGGAAGATATG-3' and 5'-  
       cccatggttaggatcctatCGATTTGGTGTATCGAGATTGG-3' as primers and plasmid pHCl (Carrer et al. 1990) as template. PCR amplification introduced an *EcoRI* site at the 5' end and *ClaI*, *BamHI* and a *NcoI* sites at the 3' end. A T

10    introduced between the *ClaI* and the *BamHI* sites eliminates an ATG and introduces an in-frame stop codon (Sriraman 2000). The Rubisco SSU transit peptides are included in *BamHI*-*NcoI* fragments. The pKO27 fragment (Pea SSU-TP22; Sequence ID No. 7) was obtained by using

15    oligonucleotides 5'-CCGGATCCAATTCAACCACAAGAACTAAC-3' and 5'-GGGGCTAGCCATGGCAGGCCACACCTGCATGCAC-3' as primers and plasmid pSSUpGEM4 as the template (Timko et al. 1985). The pKO28 fragment (Pea SSU-TP5; Sequence ID No. 6) was obtained by using oligonucleotides 5'-

20    CCGGATCCAATTCAACCACAAGAACTAAC-3' and 5'-  
       GGGGCTAGCCATGGTCAATGGGTTCAAATAGG-3' as primers and plasmid pSSUpGEM4 as the template (Timko et al. 1985). A pea SSU-TP with 23 amino acids of the mature polypeptide is shown in Sequence ID No. 8. The cre coding region

25    included in a *NcoI*-*XbaI* fragment (Sequence ID No. 3) was obtained by PCR amplification using the Cre1 5'-  
       GGGGAGCTCCATGGCTAGCTCCAATTTACT  
       GACCGTACAC-3' and Gre2 5'-GGGTCTAGACTAATCGCCATC  
       CTCGAGCAGGCGCACCATTGC-3' oligonucleotides as primers and

30    DNA isolated from *Escherichia coli* strain BNN132 (ATCC number 47059) as template. The presence of cre gene in plant nuclear DNA was confirmed by PCR amplification with the Cre 1 and Cre3 oligonucleotides. The sequence of Cre3 oligonucleotide is 5'-TCAATCGATGAGTTGCTTC-3'.

35    The *Agrobacterium nos* terminator (Tnos) is included in a

XbaI-HindIII fragment (Svab et al. 1990). The plastid targeted nuclear cre genes were introduced as EcoRI-HindIII fragments into the pPZP212 *Agrobacterium* binary vectors (Hajdukiewicz et al. 1994) to obtain plasmids pKO27 and pKO28 with twenty two and five amino acids of the mature Rubisco SSU. A schematic map of the *Agrobacterium* vectors is shown in Fig. 3.

**Transgenic plants.** Plastid transformation using the biolistic protocol, selection of transplastomic tobacco clones (RMOP medium, 500 mg/L spectinomycin dihydrochloride) and characterization of the transplastomic clones by DNA gel blot analysis was described (Svab and Maliga 1993). Transformation with *Agrobacterium* vectors pKO28 or pKO27 and regeneration of transformed tobacco plants has also been reported (Hajdukiewicz et al. 1994). Briefly, nuclear gene transformants were selected by kanamycin resistance on RMOP shoot regeneration medium containing 100 mg/L kanamycin and 500 mg/L carbenicillin. Kanamycin resistance of the shoots was confirmed by rooting on plant maintenance (RM) medium containing 100 mg/L kanamycin. Testing of 5FC cytotoxicity was carried out on RMPO medium according to published procedures (Serino and Maliga 1997).

**Transplastomic tobacco plants with a *codA* gene flanked by direct lox sites.**

Plastid transformation vector pSAC48 carries a *codA* gene in which two lox sites flank the coding region in a direct orientation. If the *codA* coding region is deleted via the lox sites, a lox site flanked by the promoter (Pr<sub>rrn</sub>) and terminator (Tr<sub>bcl</sub>) are left behind. The selective marker in pSAC48, a pPRV111B vector



derivative, is a spectinomycin resistance (*aadA*) gene (Fig. 2). Transformation with plasmid pSCAC48 yielded a number of independently transformed transplastomic lines, of which four were purified to the homoplastomic state: Nt-pSAC48-21A, Nt-pSAC48-16C, Nt-pSAC48-16CS and Nt-pSAC48-9A. These lines are considered identical other than they have been generated independently. A uniform population of transformed plastid genomes in the transplastomic plants was verified by DNA gel blot analysis (see below).

#### **Nuclear-encoded plastid-targeted Cre genes.**

To activate deletion of the plastid *>codA>* gene we introduced an engineered *cre* gene into the nucleus of the transplastomic lines encoding a plastid-targeted CRE. Targeting of nuclear-encoded plastid proteins is by an N-terminal transit peptide (TP) cleaved off during import from the cytoplasm into plastids (Soll and Tien, 1998). To ensure plastid targeting of the CRE recombinase, it was translationally fused with the Rubisco small subunit (SSU) transit peptide (Timko et al. 1985). Therefore, the product of the protein fusion is SSU-TP-CRE. Efficiency of import of chimeric proteins depends on the size of mature protein N-terminus incorporated in the construct (Wasmann et al. 1986; Lubben et al. 1989). Two chimeric *cre* genes (*Ssu-tp-cre*) were prepared, one with 5 (vector pK028) and one with 22 (plasmid pK027) amino acids of the mature SSU N-terminus, encoding SSU-TP5-CRE and SSU-TP22-CRE, respectively. These genes are also referred to as *Cre1* and *Cre2*, respectively (Table 1). The *cre* genes were expressed in the P2' promoter and Tnos terminator cassettes in the *Agrobacterium* pPZP212 binary vector which carries kanamycin resistance (*neo*) as a selectable marker (Fig. 3).

Tobacco plant transformed with *Ssu-tp5-cre* (pK037) and *Ssu-tp22-cre* (pK036) were also obtained. In these plants the nuclear *cre* is expressed from the cauliflower mosaic virus 35S promoter (Seq. ID No. 10; Timmermans et al. 1990).

Line	Plastid genotype <sup>a</sup>	Nuclear marker	
Wild-type	<i>trnV+</i> <i>aadA-</i> <i>codA-</i>		
Nt-pSAC48-21A	<i>trnV+</i> <i>aadA+</i> <i>codA+</i>		
Nt-pSAC48-16C			
Cre1-1	<i>trnV+</i> <i>aadA+</i> <i>codA-</i> <i>trnV-</i> <i>aadA-</i> <i>codA-</i>	<i>neo</i>	
Cre1-2	<i>trnV+</i> <i>aadA+</i> <i>codA-</i> <i>trnV-</i> <i>aadA-</i> <i>codA-</i>	<i>neo</i>	
Cre1-3	<i>trnV+</i> <i>aadA-</i> <i>codA-</i>	<i>neo</i>	
Cre1-4	<i>trnV-</i> <i>aadA-</i> <i>codA-</i>	<i>neo</i>	
Cre1-10	<i>trnV-</i> <i>aadA-</i> <i>codA-</i>	<i>neo</i>	
Cre2-1	<i>trnV+</i> <i>aadA+</i> <i>codA-</i>	<i>neo</i>	
Cre2-2	<i>trnV+</i> <i>aadA+</i> <i>codA-</i> <i>trnV+</i> <i>aadA*</i> <i>codA-</i> <i>trnV-</i> <i>aadA-</i> <i>codA-</i>	<i>neo</i>	
Cre2-3	<i>trnV+</i> <i>aadA+</i> <i>codA+</i> <i>trnV+</i> <i>aadA+</i> <i>codA-</i> <i>trnV+</i> <i>aadA*</i> <i>codA-</i> <i>trnV-</i> <i>aadA-</i> <i>codA-</i>	<i>neo</i>	
Cre2-4	<i>trnV+</i> <i>aadA+</i> <i>codA-</i>	<i>neo</i>	
Cre2-5	<i>trnV+</i> <i>aadA+</i> <i>codA-</i>	<i>neo</i>	
Cre2-10	<i>trnV+</i> <i>aadA+</i> <i>codA-</i> <i>trnV-</i> <i>aadA-</i> <i>codA-</i>	<i>neo</i>	
Cre1-100	<i>trnV+</i> <i>aadA-</i> <i>codA-</i>	<i>neo</i>	
Cre2-100	<i>trnV+</i> <i>aadA-</i> <i>codA-</i>	<i>neo</i>	
Cre2-200	<i>trnV+</i> <i>aadA-</i> <i>codA-</i>	<i>neo</i>	
Cre2-300	<i>trnV+</i> <i>aadA-</i> <i>codA-</i>	<i>neo</i>	

<sup>a</sup>Presence or absence of plastid gene is indicated by + or -. Since the plastid *trnV* gene is deleted in some of the lines, the wild-type plastid genotype is *trnV+* *aadA-* *codA-*.

Deletion of *codA* from the plastid genome in somatic

**cells.**

To test the efficiency of CRE-mediated deletion in somatic cells, the *Ssu-tp-cre* genes were introduced into the nucleus of the transplastomic >codA> lines by cocultivation of *Agrobacterium* and tobacco leaf disks. Plants representing 11 individual *Ssu-tp-cre* insertion events have been characterized. Five lines (Cre1-derivatives) were obtained by transformation with *Ssu-tp5-cre* gene (vector pK028) and six lines (Cre2-derivatives) were obtained by transformation with the *Ssu-tp22-cre* (vector pK027) (Table 1).

Deletion of *codA* was first tested in a DNA sample taken from one leaf of eleven kanamycin resistant shoots representing an individual integration event of the nuclear *Cre* gene. Subsequently, 4 to 7 additional leaves were sampled from six shoots to confirm that the result of the analysis is typical for the plant.

The initial DNA samples were first screened for the loss of >codA> by PCR using the 01/02 primer pair complementary to sequences in the *aadA* coding region N terminus and the *codA* promoter (Fig. 4A). Amplification with these primers yields a ~0.7-kb fragment if >codA> is deleted and a ~2.0-kb fragment if the >codA> gene is still present. Ethidium bromide stained gels of PCR products in Fig. 5 indicate complete loss of >codA> in each of the samples. A perfect, reconstituted lox site between *Prrn* and *TrbcL* was confirmed in eight clones by PCR amplification of the region with primers 01/04 from the same DNA samples and direct sequencing of the amplification product with primer 02 (not shown). In two clones (Cre1-4, Cre1-10) a fragment is missing due to deletion of *aadA* alongside with *codA* (see below).

Plastid genome structure in the initial DNA sample was determined by gel blot analysis of *ApaI*-*EcoRV* digested

total cellular DNA. The probes were the plastid targeting region and the *aadA* and *codA* coding regions. The DNA gel blots are shown in Fig. 6. The maps of the parental genomes and deletion derivatives that help to interpret these genomes are shown in Fig. 4. In the plastid tester strains expressing no CRE (Nt-pSAC48-21A, Nt-pSAC48-16C) all three probes hybridized to the same 4.9-kb DNA fragment consistent with both *codA* and *aadA* being present in all the plastid genome copies. In the SSU-TP-CRE expressing plants no 4.9-kb fragment was detectable indicating the dramatic speed by which the *>codA>* gene was eliminated from the plastid genome. CRE-mediated deletion of *>codA>* via the *lox* sites yielded the 3.6-kb fragment detected in nine of the eleven clones. The 3.6-kb fragment was the only product detected in four clones, and was present in a heteroplastomic population in five clones. Unanticipated was formation of a 1.4-kb *ApaI-EcoRV* fragment in five clones. DNA gel blot analysis confirmed that this fragment lacks both *codA* and *aadA*, and is smaller than the wild type *ApaI-EcoRV* fragment (1.9-kb). Direct sequencing of PCR products in this region confirmed deletion of *codA*, *aadA* and *trnV* by homologous recombination via the duplicated *Prrn* promoter regions. One of the *Prrn* promoters is driving *codA*, the other is upstream of the rRNA operon at its native location. Deletion of *trnV* is the reason why the *ApaI-EcoRV* fragment derived from this region (1.4-kb) is smaller than the wild-type fragment (1.9-kb). The initial DNA samples were taken from one leaf of a plant obtained by rooting the shoot obtained after transformation with the *Ssu-tp-cre* genes. To confirm that the DNA samples extracted from the leaf were typical for the plant, we have sampled several more

leaves from the same plants (Fig. 7). In four clones *codA* was excised by CRE via the *lox* sites, and the shoots were homoplastomic for the deleted genome. Two of these, Cre1-3 and Cre2-4 were further characterized by testing seven and four additional leaves of the same plants, respectively. DNA gel blot analysis of these samples confirmed a uniform deletion of >*codA*> from all genome copies. These plants are the desired final products carrying the desired plastid transgenes and lacking the undesirable selective marker. These plants and their progeny can be used directly for the production of recombinant proteins as they are free from the selectable marker gene. Furthermore, these plants are a source of engineered chloroplasts for introduction into breeding lines by sexual crossing. The seed progeny of the plants is segregating for the *Ssu-tp-cre* activator gene. Plants with the desired chloroplasts but lacking the activator gene can be identified by PCR testing for *cre* sequences. Alternatively, individuals lacking *cre* can be identified in the seed progeny by sensitivity to kanamycin, since the *Ssu-tp-cre* genes in the pK027 and pK028 *Agrobacterium* vectors are physically linked to kanamycin resistance (*neo* gene; Fig. 3). In two clones, Cre1-4 and Cre1-10, deletion of *trnV* (encoding tRNA-Val<sup>GAC</sup>), *aadA* and *codA* occurred by homologous recombination via the duplicated *Prn* promoter region. The Cre1-10 plant is homoplastomic for the deletion based on probing seven additional leaves (Fig. 7). Apparently, the one remaining *trnV* gene encoding tRNA-Val<sup>UAC</sup> is sufficient for the translation of all valine codons, or there is import of tRNA-Val<sup>GAC</sup> from the cytoplasm. In the Cre1-4 clone some of the leaves (two out of four) contained residual genome copies with *trnV* and *aadA*.

In five clones the initial DNA samples contained

more than one type of plastid genome copies. Mixed populations of plastid genome populations were confirmed in all parts of the plants by testing additional leaves (Fig. 7). Genetically stable *codA* deletion lines can be obtained from these heteroplastomic plants by testing plants regenerated from single somatic cells or individual seedlings in a segregating seed progeny.

**Deletion of *codA* from the plastid genome in the seed progeny.**

CRE-mediate deletion of the negative plastid marker *codA* in somatic cells was described in the previous section. Deletion of the plastid marker gene in the somatic cells of the transplastomic plants, without going through a sexual cycle, is highly desirable to accelerate the production of marker-free transplastomic plants. However, this approach is feasible only if there is a system for tissue culture and plant regeneration from somatic cells. Such system is unavailable for the economically important cereal crops rice and maize. As an alternative to transformation of somatic cells, we developed CRE activator lines carrying a nuclear-encoded plastid-targeted Cre to be used as the source of Cre gene when used as a pollen parent. The tobacco CRE activator lines were obtained by transforming the nucleus of wild-type plants with SSU-TP-CRE constructs. Lines in which the Cre is linked to a nuclear kanamycin resistance gene in a wild-type cytoplasm are Cre1-100, Cre2-100, Cre2-200 and Cre2-300 (Table 1).

To activate deletion of *>codA>* in the seed progeny, tester plants Nt-pSAC48-21A and Nt-pSAC48-16C were emasculated to prevent self fertilization, and fertilized with pollen from the Cre2-200 and Cre2-300 activator lines. The activator lines are primary transgenic plants ( $T_0$ ) segregating for the *Ssu-tp-cre*

gene. Therefore, a proportion of the seed progeny derived from the cross will have the activator genes while others will not. If the *codA* gene is present, the 01/02 primer pair marked in Fig. 4 amplifies a 2.0-kb fragment. If the *codA* gene is absent, the same primers will amplify a 0.7-kb fragment. PCR analysis shown in Fig. 8 confirmed CRE-mediated deletion of *>codA>* in seedlings. The Cre1-100, Cre2-100 and Cre2-300 activator lines are apparently expressing CRE efficiently, indicated by the presence of only of the 0.7-kb fragment in seedlings carrying the nuclear *cre* gene. In seedlings with no *cre* sequence the same primers amplified the 2.0-kb *codA*-containing fragment. Interestingly, *cre+* seedlings from the cross with Cre2-200 contained a mixed population of *codA* containing (2.0-kb) and *codA*-deleted (0.7-kb) fragments indicating less efficient CRE-induced deletion of *>codA>*. Thus, expression level and tissue specificity of the two nuclear *Ssu-tp22-cre* genes are characteristic for the individual transformation events. CRE activity of Cre1-100, Cre2-100 and Cre2-300 activator lines is more suitable for rapid elimination of *>codA>* in a cross than the Cre2-200 line. It is undesirable to maintain the *Ssu-tp-cre* activator genes in the production lines. However, these are encoded in the nucleus, and can be separated from the transgenic chloroplasts in the next seed progeny. Linkage of *Ssu-tp-cre* to the nuclear kanamycin resistance gene facilitates identification of seedlings lacking *cre* in a segregating seed population.

**CRE site-specific recombinase for deletion of plastid DNA sequences.** Biolistic transformation of tobacco leaves always yields shoots containing a mixed population of plastid genome copies. A mixed population

of plastid genome copies is determined by DNA gel blot analysis (Carrer et al. 1993; Svab and Maliga 1993; Carrer and Maliga 1995) and can be visualized in UV light when expressing the green fluorescence protein in plastids (Khan and Maliga 1999). Homoplastomic, genetically stable plants are obtained during a second cycle of plant regeneration from the leaves of the regenerated plants or in the seed progeny. The cells of the >codA> tester strains carry a uniform population of plastid genome copies. Thus, the *Ssu-tp-cre* is introduced into the nuclear genome of a cell that is homoplastomic for >codA>. It was expected that the regenerated shoots would contain a mixed population of plastid genome copies. Instead, all plastid genome copies lack >codA>, an evidence for the enormous selection pressure by CRE activity against plastid genome copies that carry two *lox* sites. It is important that deletion of >codA> occurs in the absence of selection against >codA> by exposure to 5-fluorocytosine. Virtually complete elimination of >codA> may also be obtained when CRE activity is introduced by crossing, using pollen of an appropriate deletion activator strain. Deletion of the selectable marker in somatic cells is the preferred choice over elimination of the marker in the seed progeny. The most important advantage is time saving. Introduction of *Ssu-tp-cre* into the nucleus of somatic cells requires only three to six weeks; *Ssu-tp-cre* segregates out in the first seed progeny. In contrast, introduction and elimination of *Ssu-tp-cre* takes one additional seed progeny, about three months.

Interestingly, genome copies with one *lox* site or no *lox* site (wild-type) are stable in CRE-expressing cells. Instability of genomes with two *lox* sites may be due to formation of linear ends during the excision



process. The linear ends may then re-circularize by homologous recombination via the *Prrn* promoter sequences yielding the *trnV-aadA-codA* deletion derivatives.

5     **CRE engineering.** Although CRE is a prokaryotic protein, it naturally carries a nuclear localization signal (NLS) that targeted a CRE-GFP fusion protein to the nucleus in mammalian cells. The NLS sequences overlap the DNA binding regions and the integrity of this region is  
10     important for DNA recombinase activity (Le et al. 1999). We targeted the newly-synthesized TP-CRE protein to plastids using a plastid-targeting transit peptide (TP). The TP is localized at the N terminus of plastid proteins and is cleaved off during import from the  
15     cytoplasm into plastids (Soll and Tien, 1998). Therefore, we translationally fused a plastid transit peptide with CRE to direct its import from the cytoplasm to plastids. Translational fusion yielded a protein with an N-terminal plastid targeting signal and an internal  
20     nuclear localization signal. Efficient CRE-mediated deletion of plastid-encoded *codA* genes indicates targeting of SSU-TP-CRE to plastids. When two potential targeting sequences are present, in general one of them out-competes the other (Small et al. 1998). N-terminal  
25     organelle targeting sequences normally dominate the second internal localization signal. For example, the 70-kDa heat shock protein of watermelon cotyledons that carry N-terminal plastidal and internal glyoxysomal targeting sequences are exclusively targeted to  
30     plastids. Proteins are localized to glyoxysomes only in the absence of the plastidal presequence (Wimmer et al. 1997). The tRNA modification enzymes contain information for both mitochondrial (N-terminal extension) and nuclear targeting. The enzyme with the N-terminal  
35     extension is targeted to mitochondria and only the short

form lacking the N-terminal extension is targeted to the nucleus (Small et al. 1998). It was fortunate, that the Rubisco SSU N-terminal transit peptide dominated the CRE nuclear localization signals and the TP-CRE fusion protein was directed to plastids (chloroplasts).

A second property that is important for the present invention is maintenance of recombinase activity when CRE is fused with proteins or peptides at its N and C termini. N-terminal fusion of CRE with the *E. coli* maltose binding protein did not interfere with recombinase function (Kolb and Siddell 1996). CRE was also shown to accept a C-terminal fusion with GFP (Le et al. 1999) as well as an 11-amino-acid epitope to the herpes simplex virus (HSV) glycoprotein D coat protein. The epitope tag facilitates detection of CRE expression *in vitro* and *in vivo* using immunofluorescent labeling with a commercially available antibody (Stricklett et al. 1998). Apparently, the five and 22 amino acids that are left behind after processing of the SSU-TP5-CRE and SU-TP22-CRE proteins did not interfere with CRE function.

**Dominant negative selection markers for positive identification of deletion derivatives.** A practical application of the present invention is the removal of selectable marker genes from the transformed plastid genome. In tobacco, the excision process mediated by the CRE constructs described herein is so efficient that the >codA< deletion derivatives can be identified in the absence of 5FC selection. However, in other crops CRE-mediated excision of marker genes may be less efficient. In these species, the positive selective marker (*aadA*) may be fused with a dominant negative selective marker using linker peptides as described in the literature (Khan and Maliga 1999) or the positive and negative

marker genes may be combined in a dicistronic operon (Staub and Maliga 1995). Dominant negative selection markers allow normally non-toxic compounds to be used as toxic agents, so that cells which express these markers are non-viable in the presence of the compound, while

For example, cytosine deaminase is absent in plants. Expression of *codA*, encoding cytosine deaminase (CD; EC 3.5.4.1), in plastids renders tissue culture cells and seedlings sensitive to 5FC, facilitating direct identification of clones lacking this negative selective marker (Serino and Maliga 1997). Cytosine deaminase converts 5-fluorocytosine (5FC) into 5-fluorouracil (5FU), the precursor of 5-fluoro-dUMP. 5FC is lethal for CD-expressing cells due to irreversible inhibition of thymidylate synthase by 5-fluoro-dUMP (Beck et al. 1972). We have found that seedlings and plant tissues expressing *>codA>* were sensitive to 5FC. Seedlings lacking *codA* could be readily identified by 5FC resistance. Thus, the constructs described here are suitable to express cytosine deaminase at sufficiently high levels to be useful to implement a negative selection scheme.

Alternative negative selective markers can be obtained by adaptation of substrate-dependent negative selection schemes described for nuclear genes. Such negative selection schemes are based on resistance to indole, naphthyl, or naphthalene acetamide (Depicker et al. 1988; Karlin-Neumann et al. 1991; Sundaresan et al. 1995), chlorate (Nussaume et al. 1991), kanamycin (Xiang and Guerra 1993) and 5-fluorocytosine (5FC) (Perera et al. 1993; Stougaard 1993).

## EXAMPLE 2

### Cre-MEDIATED INVERSION OF PLASTID DNA SEQUENCES

If the lox sites in bacteria are in an inverted orientation, CRE-mediated recombination results in an inversion of the intervening DNA. We have tested, whether the CRE-mediated inversion reaction also occurs in plastids of higher plants containing DNA sequences flanked by inverted lox sites. This was assessed using a kanamycin-resistance (>neo<) coding region in an inverted orientation relative to the promoter (Fig. 9). In this construct the non-coding strand of neo is transcribed and the plants are kanamycin sensitive. The >neo< coding region is flanked by inverted lox sites. CRE-mediated inversion of the sequences reverses neo orientation resulting in the transcription of the sense strand and expression of kanamycin resistance. Inversion of the plastid-encoded >neo< coding region may be achieved by multiple approaches. One approach is to introduce a nuclear Cre into the nucleus of somatic tobacco cells, e.g., leaf, by Agrobacterium-mediated transformation. A second approach is introduction of the nuclear-encoded Cre gene by fertilization with pollen of an appropriate activator-of-inversion strain. Additional approaches are to provide CRE-activity via the incorporation of chemically inducible promoter into the construct, or to transiently express CRE from a nuclear or chloroplast construct.

#### MATERIALS AND METHODS FOR THE PRACTICE OF EXAMPLE 2

**Plastid neo gene with inverted lox sites.** The neo gene is contained in a *SacI-HindIII* fragment. The gene map is shown in Fig. 8. PrnloxI (Seq. ID No. 1) is a plastid rRNA operon (*rrn16*) promoter derivative. It is contained in a *SacI-XbaI* fragment obtained by PCR using oligonucleotides 5'-ggggagctcGCTCCCCGCGTCGTTCAATG-3' and 5'-ggtctagataaacttcgtatagcatattatacgaagttatGCTCCC AGAAATATAGCCA-3' as primers and plasmid pZS176

(progenitor of plasmid pZS197; Svab and Maliga 1993) as a template. The promoter fragment *PrrnloxI* contains a *lox* site at the 3' end adjacent to the *XbaI* site.

The *neo* coding region is contained in an *NcoI*-*XbaI*

5 fragment derived from plasmid pHC62. The *neo* sequence in plasmid pHC62 is identical with the *neo* sequence shown in Fig. 28B, US Patent 5,877,402. The *EcoRI*-*NcoI* fragment contains the ribosome binding site from plasmid pZS176. The fragment was obtained by annealing the

10 complementary oligonucleotides 5'-  
AATTCGAAGCGCTTGGATACAGTTGTAGGGAGGGATC-3' and 5'-  
CATGGATCCCTCCCTACAACGTATCCAAGCGCTTCG-3'. The *TrbcLloxI* (Seq. ID No. 2) is the *rbcL* 3'-untranslated region contained in an *EcoRI*-*HindIII* fragment obtained by PCR

15 using oligonucleotides 5'-gggaattcataacttcgtatagcatacatatacgaagttatAGACATTAGCAGATAAATT-3' and 5'-  
gggggtaccaagcttgCTAGATTTTGTATTTCAAATCTTG-3' and plasmid pMSK48 (Khan and Maliga 1999) as template. *TrbcLloxI* contains a *lox* site adjacent to the *EcoRI* site in an

20 inverted orientation relative to the *lox* site in *PrrnloxI*. The chimeric *PrrnloxI:neo:TrbcLloxI* gene was introduced into the tobacco plastid transformation vector pPRV111B (Zoubenko et al. 1994) as a *SacI*-*HindIII* fragment to obtain plasmid pSAC38.

25

**Plastid-targeted nuclear cre linked to a nuclear gentamycin resistance (*aacC1*) gene.** The plastid targeted nuclear *cre* genes were introduced as *EcoRI*-*HindIII* fragments into the pPZP222 *Agrobacterium* binary vectors

30 which carry a plant-selectable gentamycin resistance gene (Hajdukiewicz et al. 1994) to obtain plasmids pK030 and pK031 with twenty two and five amino acids of the mature Rubisco SSU. The map of the *Agrobacterium* vectors is identical with the one shown in Fig. 3. other than

they carry a gentamycin resistance gene.

**Transplastomic tobacco plants with a neo gene flanked by inverted lox sites.**

Plastid transformation vector pSAC38 with the inverted >neo< gene is shown in Fig. 9. The inverted >neo< gene was introduced into plastids by selection for spectinomycin resistance (aadA) encoded in the vector.

Two independently transformed lines were purified to the homoplastomic state: Nt-pSAC38-9A and Nt-pSAC38-10C. The homoplastomic state was confirmed by DNA gel blot analysis.

**Nuclear-encoded plastid-targeted Cre genes.**

Plant activator lines in which *Ssu-tp-cre* is linked to a nuclear kanamycin resistance gene have been described in Example 1. The plastid marker to test CRE-activated inversion described in Example 2 utilizes a kanamycin resistance gene. Kanamycin resistance conferred by the plastid gene due to CRE-mediated inversion could not be distinguished from kanamycin resistance conferred by the marker gene of the *Agrobacterium* binary vector that was used to introduce the nuclear cre. Therefore, we have constructed activator strains in which *Ssu-tp-cre* is linked to gentamycin resistance. The *Ssu-tp22-cre* gene linked to the nuclear gentamycin resistance is the Cre3 strain and the *Ssu-tp5-cre* gene linked to gentamycin resistance is the Cre4 strain.

**Inversion of >neo< in the plastid genome of somatic cells.**

The nuclear cre genes were introduced into the

chloroplast >neo< tester strains by cocultivation of tobacco leaves with the *Agrobacterium* strains and selection for gentamycin resistance (100 mg/L). Digestion of total cellular DNA with BamHI and probing with the plastid targeting region (ApaI-EcoRV fragment, Fig. 4) hybridizes to 1.8-kb and a 3.8-kb fragments in the parental Nt-psAC38-10C lines (Fig. 10). Activation by CRE in lines Cre3-3 and Cre4-5 created a mixed population of >neo< genes representing the original and inverted orientations detected as the original 3.8-kb and 1.8-kb and the newly created 4.6-kb and 0.9-kb hybridizing fragments. Lines carrying the cre and an approximately wild-type size fragment are *aadA-neo* deletion derivatives, similar to those shown in Fig. 4. Thus, it appears that CRE mediated inversion via *lox* sites creates increased local recombination frequencies that leads to deletion of the transgenes via the short direct repeats of Prn promoters.

#### **Controlling inversion via *lox* sites by CRE activity.**

Here we describe constructs for CRE-mediated inversion of plastid genome segments flanked by inverted *lox* sites. Inversion of the sequences is independent of the encoded genetic information and relies only on CRE activity. CRE activity may be provided transiently, by expression in plastids from plastid signals described in US patent 5,877,402, or from nuclear genes encoding a plastid-targeted CRE. Such plastid-targeted CRE constructs are described in Example 1, for example the *Ssu-tp5-cre* or *Sssu-tp22-cre* genes. Alternative approaches to provide CRE activity are stable incorporation of a plastid-targeted nuclear Cre into the nucleus of somatic (leaf) cells by *Agrobacterium*-mediated, PEG induced or biolistic

transformation or by fertilization with pollen from a transformed plant. The Agrobacterium P2 promoter and cauliflower mosaic virus 35S promoter exemplified here are constitutive promoters. Regulated expression of CRE  
5 may be important for certain applications.

Developmentally timed expression may be obtained from promoters with tissue specific activity. Regulated expression of CRE may be obtained from chemically induced nuclear gene promoters responding to elicitors,  
10 steroids, copper or tetracycline (reviewed in; Gatz et al. 1992; Mett et al. 1993; Aoyama and Chau 1997; Gatz 1997; Martinez et al. 1999; Love et al. 2000) and described in US patent 5,614,395.

#### 15 **Controlled expression of deleterious gene products**

There are a variety of valuable heterologous proteins that interfere with plastid metabolism. For example, certain proteins may be inserted into photosynthetic membranes and interfere with  
20 photosynthesis. This problem can be circumvented by first growing the plants to maturity, then activating production of the deleterious protein by chemically inducing CRE expression. CRE, in turn, will make the gene expressible by lox-mediated inversion of the coding  
25 region.

The molecular tools necessary for the construction of such plastid genes are described in present application. In case of the monocistronic inversion vector the gene of interest (*goi*) is flanked by inverted  
30 lox sites and is introduced by linkage with *aadA* (Fig. 12). The selectable marker (*aadA*) coding region is the first reading frame, and is expressed from the promoter. The *goi* reading frame is the second coding region, and it is not expressed as it is in an inverted orientation  
35 relative to the promoter. Expression of *goi* is induced



by CRE-mediated inversion of the *goi* coding region, as described for >neo< in Example 2 and is shown in Fig. 12.

5 The dicistronic lox inversion vector is shown in Fig. 13. In this case the inverted lox sites flank both *aadA* and *goi*. The selectable marker (*aadA*) coding region is expressed from the promoter. The *goi* reading frame is not expressed as it is in an inverted orientation relative to the promoter. Expression of *goi* is induced  
10 by CRE-mediated inversion of the *aadA*-*goi* containing region that results in simultaneous expression of *goi* and inactivation of *aadA*.

The presence of two lox sites may destabilize the plastid genome that leads to CRE-independent deletion of  
15 plastid genome sequences. However, it appears that CRE activity by itself is not mutagenic, and the plastid genomes are stable if only one lox site is present. Mutant lox sites that are efficiently excised but recombine into excision resistant sites have been  
20 described (Hoess et al. 1982; Albert et al. 1995). Such lox sites would mediate efficient inversion, but the new lox sites would be resistant to additional cycles of CRE activation. Providing only a short burst of CRE activation using a chemically induced promoter  
25 could further refine the expression system.

### EXAMPLE 3

#### CRE-MEDIATED DELETION TO OBTAIN MARKER FREE TRANSPLASTOMIC PLANTS AND FOR HIGH LEVEL EXPRESSION OF 30 THE RECOMBINANT PROTEINS

Plastid *loxP* vectors in this section are described for CRE-mediated excision of selective markers in transplastomic plants. Since excision of sequences

between directly oriented *lox* sites is very efficient, variants of the same vectors can be used for CRE-activated expression of recombinant proteins. A family of plastid vectors with suitably positioned *lox* sites is shown schematically in Fig. 14 through Fig. 17.

The map of the basic tobacco plastid *lox* deletion vector is shown in Fig. 14. It contains (a) two directly oriented *lox* sites separated by a unique *Bgl*III cloning site and (b) an adjacent polycloning site. These sequences (Seg. ID No. 11) are inserted into the *Sca*I site plastid repeat vector pPRV100 (US Patent 5,877,402; Zoubenko et al. 1994). Suitable marker genes (*aadA*, *neo* or *kan*, *bar*, glyphosate resistance, bromoxynil resistance) for insertion into the *Bgl*III site have been described in US Patent 5,877,402, WO 00/07421 and WO 00/03022.

The map of the tobacco plastid *lox* >*aadA*> deletion vector is shown in Fig. 15. It is the basic *lox* deletion vector with an *aadA* gene cloned into the *Bgl*III sites oriented towards the *rrn* operon.

Maps of constitutive *lox* dicistronic deletion vectors are shown in Fig. 16 through Fig. 18. This dicistronic design enables simultaneous expression of both the first and the second open reading frames. The selectable marker designed for excision may be encoded in the first (Fig. 16) or second (Fig. 17, Fig. 18) open reading frames. Since a minimally 34 bp *lox* site is located between the two reading frames, both the marker gene (*aadA*) and the gene of interest have their own leader sequence to facilitate translation (Fig. 16, Fig. 17). Translational coupling may also be feasible if the *lox* site is incorporated in the marker gene coding region N terminus (Fig. 18). DNA sequence of promoter-*lox* constructs shown in Figs. 16 is set forth in Seq. ID

No. 1. Promoters and promoter-leader combinations suitable to promote high-level protein expression in plastids are described in European Patent Applications WO 00/07421, WO 97/06250 and WO 98/55595. Sequences suitable for directly oriented *lox* sites are given in Seq. ID No. 11. Translational coupling between a gene of interest and the downstream *aadA* is shown in Fig. 18. There are multiple ways of achieving translational coupling between adjacent genes (Baneyx 1999). One approach is incorporation of a properly spaced ribosome binding-site in the upstream gene's coding region (Schoner et al. 1986; Omer et al. 1995). An example for a suitable sequence directly upstream of the translation initiation codon (ATG ) would be G-GAG-GAA-TAA-CTT-ATG. A specific example for the use of the sequence is translational coupling between a *bar* (suitable source described in European Patent Application WO 00/07421) and a downstream *aadA* are given in Seq. ID No. 12. Note *SalI* site downstream of AUG incorporated to facilitate engineering the *BglII-SalI* region and the directly oriented *lox* sites in the *aadA* coding region and downstream of *aadA*. The sequence is given for a *BglII-SpeI* fragment. The *BglII* site is within the *bar* coding region; the *SpeI* site is downstream of the second *lox* site, as marked in Fig. 18. If a C-terminal extension to create a ribosome binding site is unacceptable, a suitable sequence may be obtained by silent mutagenesis of the coding region at the third codon position. Variants of plastid ribosome binding sites have been catalogued (Bonham-Smith and Bourque 1989)

A tobacco inducible *lox* deletion vector is shown in Fig. 19. The marker gene (*aadA*) is encoded in the first reading frame, followed by a silent *goi* lacking the translation initiation codon (ATG) and the 5'

untranslated leader. Expression of the *goi* frame is triggered by *aadA* excision that results in translational fusion of the *aadA* N-terminal region with the *goi*. After *aadA* excision the *goi* mRNA is translated from the *aadA* translation control signals, the 5' UTR and AUG. DNA sequence of the SacI-NheI fragment is given in Seq. ID. No. 13. The *Prn* promoter-*atpB* translational control region is described in European Patent Application WO 00/07421. The *aadA* construct has two directly-oriented *lox* sites: one in the coding region N-terminus and one downstream of *aadA* to facilitate CRE-mediated excision of the marker gene.

#### EXAMPLE 4

##### DELETION OF VITAL PLASTID GENES TO OBTAIN CYTOPLASMIC MALE STERILITY

US Patent 5,530,191 provides a cytoplasmic male sterility (CMS) system for plants, which is based on modification of the plastid genome. The CMS system comprises three transgenes: a "plastid male sterility" gene that causes plastid and cellular disablement of the anther tissue, and two nuclear genes that regulate the expression of the plastid gene. An important feature of the system is developmentally timed cellular death based on the expression, or the lack of the expression, of a plastid gene. As one specific approach to induce developmentally timed ablation of anther tissue we describe CRE-mediate excision of essential plastid genes via directly oriented *lox* sites.

The number of genes encoded by the plastid genome is about 120. Some of the genes are non-essential and may be inactivated by targeted gene disruption without a major phenotypic consequence. Good examples are the plastid *ndh* genes (Burrows et al. 1998; Shikanai et al.

1998) or the *trnV* gene the deletion of which has been described in Example 1. Excision of these genes is unlikely to cause cell ablation. The photosynthetic genes are essential for survival under field conditions. However, pigment deficient, non-photosynthetic plants can be maintained as long as they are grown on a sucrose-containing medium, or are grafted onto photosynthetically active wild-type (green) plants (Kanevski and Maliga 1994). Some of the house-keeping genes, such as the genes encoding the plastid multisubunit RNA polymerase are essential for photosynthetic growth, but not for survival (Allison et al. 1996). Thus, deletion of these genes is not suitable to trigger cell death. Only a relatively small number of plastid genes have proven to be essential for viability. The essential nature of the genes was recognized by the lack of homoplastomic cells in gene disruption experiments indicating that the loss of these genes results in cellular death. Cellular death due to lack of plastid function is understandable, as plastids are the site of the biosynthesis of amino acids, several lipids and are required for nitrate assimilation. Examples of plastid genes essential for cellular survival are the *clpP* protease subunit gene (Huang et al. 1994), *ycf1* and *ycf2*, the two largest plastid-encoded open reading frames (Drescher et al. 2000).

To induce cellular death by CRE-mediated excision, directly oriented *lox* sites can be incorporated in the plastid genome flanking essential genes, as shown for *clpP* in Fig. 20. The *clpP* gene has two large introns (807 bp and 637 bp) and the region can be conveniently cloned as a *SalI*-*SphI* fragment. The selectable marker *aadA* is inserted into a *KpnI* restriction site created by PCR mutagenesis downstream of *clpP* Exon 3, oriented towards *rps12* Exon I. One of the *lox* sites is

engineered next to the *aadA* gene, the second *lox* site is inserted in Intron I. Cellular death is induced by activation of the nuclear *Cre* gene as described in US Patent 5,530,191. It is necessary to use a selective  
 5 marker, such as *aadA* to introduce the *lox* sites into the plastid genome. The *aadA* gene can subsequently eliminated using a second, independent site specific recombinase such as FRT via the *frt* sites engineered into the transformation vector shown in Fig. 20.

10 Alternative targets for CRE-mediated deletion in a CMS system are the essential ribosomal protein genes such as *rpl23*, the ribosomal RNA operon (for insertion sites see; Staub and Maliga 1992; Zoubenko et al. 1994) and the *ycf1* and *ycf2* genes (Drescher et al. 2000)

15 The following sequences are referred to throughout the specification and facilitate the practice of the present invention.

SEQ. No. 1: *PrnloxI*. sequence

20 gagctcGCTCCCCGCGTCGTTCAATGAGAAATGGATAAGAGGCTCGTGGGATTGA  
 CGTGAGGGGGCAGGGATGGCTATATTTCTGGGAGCataacttcgtataatgtatgc  
 tatacgaagttatctaga

25 SEQ. No. 2: *TrbcLloxI*. sequence

gaattcataacttcgtatagcatatacattatacgaagttatAGACATTAGCAGATAA  
 ATTAGCAGGAAATAAAGAAGGATAAGGAGAAAGAACTCAAGTAATTATCCTTCGTT  
 CTCTTAATTGAATTGCAATTAAACTCGGCCCAATCTTTTACTAAAAGGATTGAGCC  
 30 GAATACAACAAAGATTCTATTGCATATATTTTGACTAAGTATATACTTACCTAGAT  
 ATACAAGATTTGAAATACAAAATCTAGcaagcttggtacc

SEQ. No. 3: *cre* coding region. sequence

35 gagctccATGgctagcTCC AATTTACTGA CCGTACACCA AAATTTGCCT  
 GCATTACCGG TCGATGCAAC GAGTGATGAG GTTCGCAAGA ACCTGATGGA  
 CATGTTCCAGG GATCGCCAGG CGTTTTCTGA GCATACCTGG AAAATGCTTC  
 TGTCCGTTTG CCGGTCGTGG GCGGCATGGT GCAAGTTGAA TAACCGGAAA  
 40 TGGTTTCCCG CAGAACCTGA AGATGTTTCG GATTATCTTC TATATCTTCA  
 GGCGCGCGGT CTGGCAGTAA AAATATCCA GCAACATTG GGCCAGCTAA  
 ACATGCTTCA TCGTCGGTCC GGGCTGCCAC GACCAAGTGA CAGCAATGCT  
 GTTTCACCTGG TTATGCGGCG GATCCGAAAA GAAAACGTTG ATGCCGGTGA

5 ACGTGCAAAA CAGGCTCTAG CGTTCGAACG CACTGATTTC GACCAGGTTC  
 GTTCACTCAT GGAAAATAGC GATCGCTGCC AGGATATACG TAATCTGGCA  
 TTTCTGGGGA TTGCTTATAA CACCCTGTTA CGTATAGCCG AAATTGCCAG  
 GATCAGGGTT AAAGATATCT CACGTACTGA CGGTGGGAGA ATGTTAATCC  
 10 ATATTGGCAG AACGAAAACG CTGGTTAGCA CCGCAGGTGT AGAGAAGGCA  
 CTTAGCCTGG GGGTAACTAA ACTGGTCGAG CGATGGATTT CCGTCTCTGG  
 TGTAGCTGAT GATCCGAATA ACTACCTGTT TTGCCGGGTC AGAAAAAATG  
 GTGTTGCCGC GCCATCTGCC ACCAGCCAGC TATCAACTCG CGCCCTGGAA  
 GGGATTTTTG AAGCAACTCA TCGATTGATT TACGGCGCTA AGGATGACTC  
 15 TGGTCAGAGA TACCTGGCCT GGTCTGGACA CAGTGCCCGT GTCGGAGCCG  
 CGCGAGATAT GGCCCGCGCT GGAGTTTCAA TACCGGAGAT CATGCAAGCT  
 GGTGGCTGGA CCAATGTAAA TATTGTCATG AACTATATCC GTAACCTGGA  
 TAGTGAAACA GGGGCAATGG TGCGCCTGCT cGAgGATGGC GATTAGtctaga

15

SEQ. No. 4: PrnrloxD. Sequence

gagctcGCTCCCCCGCCGTCGTTCAATGAGAATGGATAAGAGGCTCGTGCGGATTGA  
 20 CGTGAGGGGGCAGGGATGGCTATATTTCTGGGAGCataacttcgtataatgtatgc  
 tatacgaagttatgaattc

SEQ. No. 5: TrbcLloxD. sequence

25 tctagataacttcgtataatgtatgctatacgaagttatAGACATTAGCAGATAAAA  
 TTAGCAGGAAATAAAGAAGGATAAGGAGAAAGAACTCAAGTAATTATCCTTCGTTT  
 TCTTAATTGAATTGCAATTAAACTCGGCCCAATCTTTTACTAAAAGGATTGAGCCG  
 AATACAACAAAGATTCTATTGCATATATTTTACTAAGTATATACTTACCTAGATA  
 TACAAGATTTGAAATACAAATCTAGcaagcttggtacc

30

SEQ. No. 6: Pea ssuTP5. sequence

35 ccggatccAA TTCAACCACA AGAACTAACA AAGTCAGAAA AATGGCTTCT  
 ATGATATCCT CTTCCGCTGT GACAACAGTC AGCCGTGCTT CTAGGGTGCA  
 ATCCGCGGCA GTGGCTCCAT TCGGCGGCCT GAAATCCATG ACTGGATTCC  
 CAGTGAAGAA GGTCAACACT GACATTACTT CCATTACAAG CAATGGTGGA  
 AGAGTAAAGT GCATGCAGGT GTGGCCTgcc atggctagc

40

SEQ. No. 7: Pea ssuTP22. sequence

45 ccggatcc AA TTCAACCACA AGAACTAACA AAGTCAGAAA AATGGCTTCT  
 ATGATATCCT CTTCCGCTGT GACAACAGTC AGCCGTGCTT CTAGGGTGCA  
 ATCCGCGGCA GTGGCTCCAT TCGGCGGCCT GAAATCCATG ACTGGATTCC  
 CAGTGAAGAA GGTCAACACT GACATTACTT CCATTACAAG CAATGGTGGA  
 AGAGTAAAGT GCATGCAGGT GTGGCCTCCA ATTGGAAAGA AGAAGTTTGA  
 GACTCTTTCC TATTTGCCAC CATTGACCat ggctagc

50

SEQ. No. 8: Pea ssuTP23. sequence

ccggatccAA TTCAACCACA AGAACTAACA AAGTCAGAAA AATGGCTTCT  
 ATGATATCCT CTTCCGCTGT GACAACAGTC AGCCGTGCTT CTAGGGTGCA  
 ATCCGCGGCA GTGGCTCCAT TCGGCGGCCT GAAATCCATG ACTGGATTCC  
 CAGTGAAGAA GGTCAACACT GACATTACTT CCATTACAAG CAATGGTGGA  
 5 AGAGTAAAGT GCATGCAGGT GTGGCCTCCA ATTGGAAAGA AGAAGTTTGA  
 GACTCTTTCC TATTTGCCAC CATTGACCAG AGATCAGTTG gctagcgg

SEQ. No. 9: P2 promoter sequence

10 gaattCATTT TCACGTGTGG AAGATATGAA TTTTTTTGAG AAAC TAGATA  
 AGATTAATGA ATATCGGTGT TTTGGTTTTT TCTTGTGGCC GTCTTTGTTT  
 ATATTGAGAT TTTTCAAATC AGTGCGCAAG ACGTGACGTA AGTATCTGAG  
 CTAGTTTTTA TTTTCTACT AATTTGGTCG TTTATTTTCGG CGTGTAGGAC  
 ATGGCAACCG GGCCTGAATT TCGCGGGTAT TCTGTTTCTA TTCCAAC TTT  
 15 TTCTTGATCC GCAGCCATTA ACGACTTTTG AATAGATACG CTGACACGCC  
 AAGCCTCGCT AGTCAAAAGT GTACCAAACA ACGCTTTACA GCAAGAACGG  
 AATGCGCGTG ACGCTCGCGG TGACGCCATT TCGCCTTTTC AGAAATGGAT  
 AAATAGCCTT GCTTCCTATT ATATCTTCCC AAATTACCAA TACATTACAC  
 TAGCATCTGA ATTCATAAC CAATCTCGAT ACACCAAATC GATaggatcc  
 20 taccatgg

SEQ. No. 10: 35S promoter sequence

AAGCTTGCCA ACATGGTGGA GCACGACACT CTCGTCTACT CCAAGAATAT  
 25 CAAAGATACA GTCTCAGAAG ACCAAAGGGC TATTGAGACT TTTCAACAAA  
 GGGTAATATC GGGAAACCTC CTCGATTCC ATTGCCCAGC TATCTGTCAC  
 TTCATCAAAA GGACAGTAGA AAAGGAAGGT GGCACCTACA AATGCCATCA  
 TTGCGATAAA GGAAAGGCTA TCGTTCAAGA TGCCTCTGCC GACAGTGGTC  
 CCAAAGATGG ACCCCCACCC ACGAGGAGCA TCGTGGA AAA AGAAGACGTT  
 30 CCAACCACGT CTTCAAAGCA AGTGGATTGA TGTGATAACA TGGTGGAGCA  
 CGACACTCTC GTCTACTCCA AGAATATCAA AGATACAGTC TCAGAAGACC  
 AAAGGGCTAT TGAGACTTTT CAACAAAGGG TAATATCGGG AAACCTCCTC  
 GGATTCCATT GCCCAGCTAT CTGTCACTTC ATCAAAAGGA CAGTAGAAAA  
 GGAAGGTGGC ACCTACAAAT GCCATCATTG CGATAAAGGA AAGGCTATCG  
 35 TTCAAGATGC CTCTGCCGAC AGTGGTCCCA AAGATGGACC CCCACCCACG  
 AGGAGCATCG TGGAAAAAGA AGACGTTC CAACAGTCTT CAAAGCAAGT  
 GGATTGATGT GATATCTCCA CTGACGTAAG GGATGACGCA CAATCCCACT  
 ATCCTTCGCA AGACCCTTCC TCTATATAAG GAAGTTCATT TCATTTGGAG  
 AGGACACGCT GAAATCACCA GTCTCTCTCT ACAAATCTAT CTCTCTCGAT  
 40 TCGCGAGCTC GGTACCCGGG gatcgatcc

SEQ. No. 11: KpnI-lox-BglIII-lox-HindIII fragment

45 ggtaccATAACTTCGTATAATGTATGCTATACGAAGTTATagatctATAACTTCGT  
 ATAATGTATGCTATACGAAGTTATAagctt

Seq. ID No. 12. Translational coupling of bar and aadA according to scheme in Fig. 18. BglIII-SpeI fragment.

50 GAGATCTGgg aggaataact tATGgggggtc gacATAACTT CGTATAATGT  
 ATGCTATACG AAGTTATtaG AAGCGGTGAT CGCCGAAGTA TCGACTCAAC



5 TATCAGAGGT AGTTGGCGTC ATCGAGCGCC ATCTCGAACC GACGTTGCTG  
 GCCGTACATT TGTACGGCTC CGCAGTGGAT GGCGGCCTGA AGCCACACAG  
 TGATATTGAT TTGCTGGTTA CCGTGACCGT AAGGCTTGAT GAAACAACGC  
 GGCGAGCTTT GATCAACGAC CTTTTGGAAA CTTCCGGCTTC CCCTGGAGAG  
 10 AGCGAGATTC TCCGCGCTGT AGAAGTCACC ATTGTTGTGC ACGACGACAT  
 CATTCGCTGG CGTTATCCAG CTAAGCGCGA ACTGCAATTT GGAGAATGGC  
 AGCGCAATGA CATTCCTTGCA GGTATCTTCG AGCCAGCCAC GATCGACATT  
 GATCTGGCTA TCTTGCTGAC AAAAGCAAGA GAACATAGCG TTGCCTTGGT  
 AGGTCCAGCG GCGGAGGAAC TCTTTGATCC GGTTCCTGAA CAGGATCTAT  
 15 TTGAGGCGCT AAATGAAACC TTAACGCTAT GGAACCTCGC GCCCGACTGG  
 GCTGGCGATG AGCGAAATGT AGTGCTTACG TTGTCCCGCA TTTGGTACAG  
 CGCAGTAACC GGCAAAATCG CGCCGAAGGA TGTCGCTGCC GACTGGGCAA  
 TGGAGCGCCT GCCGGCCCAG TATCAGCCCG TCATACTTGA AGCTAGACAG  
 GCTTATCTTG GACAAGAAGA AGATCGCTTG GCCTCGCGCG CAGATCAGTT  
 20 GGAAGAATTT GTCCACTACG TGAAAGGCGA GATCACCAG GTAGTCGGCA  
 AATAAATAAC TTCGTATAAT GTATGCTATA CGAAGTTaAa ctagt

Seq. ID No. 13. CRE-induced expression of recombinant  
 protein according to design in Fig. 19. SacI-NheI  
 fragment.

25 gagctcGCTC CCCC GCCGTC GTTCAATGAG AATGGATAAG AGGCTCGTGG  
 GATTGACGTG AGGGGGCAGG GATGGCTATA TTTCTGGGAG AATTAACCGA  
 TCGACGTGCa AGCGGACATT TATTTTaAAT TCGATAATTT TTGCAAAAAC  
 ATTTTCGACAT ATTTATTTAT TTTATTATTA TGgggATAAC TTCGTATAAT  
 GTATGCTATA CGAAGTTATt aGAAGCGGTG ATCGCCGAAG TATCGACTCA  
 ACTATCAGAG GTAGTTGGCG TCATCGAGCG CCATCTCGAA CCGACGTTGC  
 TGGCCGTACA TTTGTACGGC TCCGAGTGG ATGGCGGCCT GAAGCCACAC  
 30 AGTGATATTG ATTTGCTGGT TACGGTGACC GTAAGGCTTG ATGAAACAAC  
 GCGGCGAGCT TTGATCAACG ACCTTTTGGA AACTTCGGCT TCCCCTGGAG  
 AGAGCGAGAT TCTCCGCGCT GTAGAAGTCA CCATTGTTGT GCACGACGAC  
 ATCATTCCTG GCGGTTATCC AGCTAAGCGC GAACTGCAAT TTGGAGAATG  
 GCAGCGCAAT GACATTCTTG CAGGTATCTT CGAGCCAGCC ACGATCGACA  
 TTGATCTGGC TATCTTGCTG ACAAAGCAA GAGAACATAG CGTTGCCTTG  
 35 GTAGGTCCAG CGGCGGAGGA ACTCTTTGAT CCGGTTCCCTG AACAGGATCT  
 ATTTGAGGCG CTAAATGAAA CCTTAACGCT ATGGAACCTG CCGCCCGACT  
 GGGCTGGCGA TGAGCGAAAT GTAGTGCTTA CGTTGTCCCG CATTTGGTAC  
 AGCGCAGTAA CCGGCAAAAT CGCGCCGAAG GATGTGCTG CCGACTGGGC  
 AATGGAGCGC CTGCCGGCCC AGTATCAGCC CGTCATACTT GAAGCTAGAC  
 40 AGGCTTATCT TGGACAAGAA GAAGATCGCT TGGCCTCGCG CGCAGATCAG  
 TTGGAAGAAT TTGTCCAATA CGTGAAAGGC GAGATCACCAG AGGTAGTCGG  
 CAAATAAATA ACTTCGTATA ATGTATGCTA TACGAAGTTA Ttagctagc

45

#### REFERENCES

Adams DE, Bliska JB, Cozzarelli NR (1992) Cre-lox  
 recombination in Escherichia coli cells. Mechanistic  
 differences from the in vitro reaction. J Mol Biol  
 226:661-673

50

- Albert H, Dale EC, Lee E, Ow DW (1995) Site-specific integration of DNA into wild-type and mutant lox sites placed in the plant genome. *Plant J* 7:649-659
- 5 Allison LA, Simon LD, Maliga P (1996) Deletion of *rpoB* reveals a second distinct transcription system in plastids of higher plants. *EMBO J* 15:2802-2809
- 10 Aoyama T, Chau NH (1997) A glucocorticoid-mediated transcriptional induction system in transgenic plants. *Plant J* 11:605-612
- 15 Baneyx F (1999) Recombinant protein expression in *Escherichia coli*. *Curr Opin Biotechnol* 10:411-421
- Beck CF, Ingraham JL, Neuhard J, Thomassen E (1972) Metabolism of pyrimidines and pyrimidine nucleosides by *Salmonella typhimurium*. *J Bacteriol* 110:219-228
- 20 Bonham-Smith PC, Bourque DP (1989) Translation of chloroplast-encoded mRNA: potential initiation and termination signals. *Nucleic Acids Res* 17:2057-2080
- 25 Burrows PA, Sazanov LA, Svab Z, Maliga P, Nixon PJ (1998) Identification of a functional respiratory complex in chloroplasts through analysis of tobacco mutants containing disrupted plastid *ndh* genes. *EMBO J* 17:868-876
- 30 Carrer H, Hockenberry TN, Svab Z, Maliga P (1993) Kanamycin resistance as a selectable marker for plastid transformation in tobacco. *Mol Gen Genet* 241:49-56
- 35 Carrer H, Maliga P (1995) Targeted insertion of foreign genes into the tobacco plastid genome without physical linkage to the selectable marker gene. *Biotechnology* 13:791-794
- 40 Carrer H, Staub JM, Maliga P (1990) Gentamycin resistance in *Nicotiana* conferred by AAC(3)-I, a narrow substrate specificity acetyl transferase. *Plant Mol Biol* 17:301-303
- 45 Craig NL (1988) The mechanism of conservative site-specific recombination. *Annual Review Of Genetics* 22:77-105
- 50 Dale EC, Ow DW (1991) Gene transfer with subsequent removal of the selection gene from the host genome. *Proc Natl Acad Sci USA* 88:10558-10562

- Depicker AG, Jacobs AM, Montagu MC (1988) A negative selection scheme for tobacco protoplast-derived cells expressing the T-DNA gene 2. *Plant Cell Rep* 7:63-66
- 5     Drescher A, Ruf S, Calsa T, Carrer H, Bock R (2000) The two largest chloroplast genome-encoded open reading frames of higher plants are essential genes. *The Plant Journal* 22:97-104
- 10    Dršge M, Pöhler A, Selbitschka W (1998) Horizontal gene transfer as a biosafety issue: a natural phenomenon of public concern. *J Biotechnol* 64:75-90
- 15    Gatz C (1997) Chemical control of gene expression. *Ann Rev Plant Physiol Plant Mol Biol* 48:89-108
- 20    Gatz C, Froberg C, Wendenburg R (1992) Stringent repression and homogeneous de-repression by tetracycline of a modified CaMV 35S promoter in intact transgenic tobacco plants. *Plant J* 2:397-404
- 25    Hajdukiewicz P, Svab Z, Maliga P (1994) The small, versatile pPZP family of *Agrobacterium* binary vectors for plant transformation. *Plant Mol Biol* 25:989-994
- 30    Hoess RH, Ziese M, Sternberg N (1982) P1 site-specific recombination: nucleotide sequence of the recombining sites. *Proc Natl Acad Sci USA* 79:3398-3402
- 35    Huang C, Wang S, Chen L, Lemieux C, Otis C, Turmel M, Liu XQ (1994) The *Chlamydomonas* chloroplast *clpP* gene contains translated large insertion sequences and is essential for cell growth. *Molecular and Genetal Genetics* 244:151-159
- 40    Kanevski I, Maliga P (1994) Relocation of the plastid *rbcL* gene to the nucleus yields functional ribulose-1,5-bisphosphate carboxylase in tobacco chloroplasts. *Proc Natl Acad Sci USA* 91:1969-1973
- 45    Karlin-Neumann GA, Brusslan JA, Tobin EM (1991) Phytochrome control of the *tms2* gene in transgenic *Arabidopsis*: a strategy for selecting mutants in the signal transduction pathway. *Plant Cell* 3:573-582
- 50    Khan MS, Maliga P (1999) Fluorescent antibiotic resistance marker to track plastid transformation in higher plants. *Nat Biotechnol* 17:910-915
- 50    Kolb AF, Siddell SG (1996) Genomic targeting with an MBP-Cre fusion protein [published erratum appears in *Gene* 1997 Apr 11;189(1):149]. *gene* 183:53-60

- Le Y, Gagneten S, Tombaccini D, Bethke B, Sauer B (1999) Nuclear targeting determinants of the phage P1 Cre DNA recombinase. *Nucleic Acids Res* 27:4703-4709
- 5 Lichtenstein C, Barrena E (1993) Prospects for reverse genetics in plants using recombination [news]. *Plant Mol Biol* 21:v-xii
- 10 Love J, Scott AC, Thompson WF (2000) Stringent control of transgene expression in *Arabidopsis thaliana* using the Top10 promoter system. *Plant J* 21:579-588
- 15 Lubben TH, Gatenby AA, Ahlquist P, Keegstra K (1989) Chloroplast import characteristics of chimeric proteins. *Plant Mol Biol* 12:13-18
- 20 Lyznik LA, Hirayama L, Rao KV, Abad A, Hodges TK (1995) Heat-inducible expression of FLP gene in maize cells. *Plant J* 8:177-186
- 25 Lyznik LA, Mitchell JC, Hirayama L, Hodges TK (1993) Activity of yeast FLP recombinase in maize and rice protoplasts. *Nucleic Acids Res* 21:969-975
- 30 Maliga P (1993) Towards plastid transformation in higher plants. *Trends Biotech* 11:101-107
- 35 Martinez A, Sparks C, Hart CA, Thompson J, Jepson I (1999) Ecdysone agonist inducible transcription in transgenic tobacco plants. *Plant J* 19:97-106
- 40 Mett VL, Lochhead LP, Reynolds PHS (1993) Copper-controllable gene expression for whole plants. *Proc Natl Acad Sci USA* 90:4567-4571
- 45 Morris AC, Schaub TL, James AA (1991) FLP-mediated recombination in the vector mosquito, *Aedes aegypti*. *Nucleic Acids Res* 19:5895-5900
- 50 Nussaume L, Vincentz M, Caboche M (1991) Constitutive nitrate reductase: a dominant conditional marker for plant genetics. *Plant J* 1:267-274
- O'Gorman S, Fox DT, Wahl GM (1991) Recombinase-mediated gene activation and site-specific integration in mammalian cells. *Science* 251:1351-1355

- Omer CA, Diehl RE, Kral AM (1995) Bacterial expression and purification of human protein prenyltransferases using epitope-tagged, translationally coupled systems. *Meth Enzymol* 250:3-12
- 5 Perera RJ, Linard CG, Signer ER (1993) Cytosine deaminase as a negative selective marker for *Arabidopsis*. *Plant Mol Biol* 23:793-799
- 10 Russell SH, Hoopes JL, Odell JT (1992) Directed excision of a transgene from the plant genome. *Mol Gen Genet* 234:49-59
- 15 Schoner BE, Belagaje RM, Schoner RG (1986) Translation of a synthetic two-cistron mRNA in *Escherichia coli*. *Proc Natl Acad Sci USA* 83:8506-8510
- 20 Serino G, Maliga P (1997) A negative selection scheme based on the expression of cytosine deaminase in plastids. *Plant J* 12:697-701
- 25 Shikanai T, Endo T, Hashimoto T, Yamada Y, Asada K, Yokota A (1998) Directed disruption of the tobacco *ndhB* gene impairs cyclic electron flow around photosystem I. *Proc Natl Acad Sci USA* 95:9705-9709
- 30 Shinozaki K, Ohme M, Tanaka M, Wakasugi T, Hayashida N, Matsabayashi T, Zaita N, Chungwongse J, Obokata J, Yamaguchi-Shinozaki K, Deno H, Kamogashira T, Yamada K, Kasuda J, Takaiwa F, Kato A, Todoh N, Shimada H, Sugiura M (1986) The complete sequence of the tobacco chloroplast genome: its gene organization and expression. *EMBO J* 5:2043-2049
- 35 Small I, Wintz H, Akashi K, Mireau H (1998) Two birds with one stone: genes that encode products targeted to two or more compartments. *Plant Mol Biol* 38:265-277
- 40 Soll J, Tien R (1998) Protein translocation into and across the chloroplastic envelope membranes. *Plant Mol Biol* 38:191-207
- 45 Sriraman P (2000) Identification and characterization of components of the plastid transcription machinery. Identification and characterization of components of the plastid transcription machinery. Rutgers University, Piscataway, NJ
- 50 Srivastava V, Anderson OD, Ow DW (1999) Single-copy transgenic wheat generated through the resolution of complex integration patterns. *Proceedings of the National Academy of Sciences* 96:11117-11121

- Staub JM, Maliga P (1992) Long regions of homologous DNA are incorporated into the tobacco plastid genome by transformation. *Plant Cell* 4:39-45
- 5 Staub JM, Maliga P (1995) Expression of a chimeric uidA gene indicates that polycistronic mRNAs are efficiently translated in tobacco plastids. *Plant J* 7:845-848
- 10 Stougaard J (1993) Substrate-dependent negative selection in plants using a bacterial cytosine deaminase gene. *Plant J* 3:755-761
- 15 Stricklett PK, Nelson RD, Kohan DE (1998) Site-specific recombination using an epitope tagged bacteriophage P1 Cre recombinase. *gene* 215:415-423
- 20 Sundaresan V, Springer P, Volpe T, Haward S, Jones JD, Dean C, Ma H, Martienssen R (1995) Patterns of gene action in plant development revealed by enhancer trap and gene trap transposable elements. *Genes Dev* 9:1797-1810
- 25 Svab Z, Harper EC, Jones JD, Maliga P (1990) Aminoglycoside-3''-adenyltransferase confers resistance to spectinomycin and streptomycin in *Nicotiana tabacum*. *Plant Mol Biol* 14:197-205
- 30 Svab Z, Maliga P (1993) High-frequency plastid transformation in tobacco by selection for a chimeric *aadA* gene. *Proc Natl Acad Sci USA* 90:913-917
- Sylvanen M (1999) In search of horizontal gene transfer. *Nat Biotechnol* 17:833
- 35 Tepfer D (1989) Ri T-DNA from *Agrobacterium rhizogenes*: a source of genes having applications in rhizosphere biology and plant development, ecology and evolution. In: Kosuge T, Nester EW (eds) *Plant-Microbe Interactions. Molecular and Genetic Perspectives*, Vol. 3, McGraw-Hill, New York, pp 294-342
- 40 Timko MP, Kaush AP, Hand JM, Cashmore AR (1985) Structure and expression of nuclear genes encoding polypeptides of the photosynthetic apparatus. In: Steinback KE, Bonitz S, Arntzen CJ, Bogorad L (eds) *Molecular biology of the photosynthetic apparatus.*, Cold Spring Harbor Laboratory, Cold Spring Harbor, pp 381-396.
- 45 Timmermans MCP, Maliga P, Vieira J, Messing J (1990) The pFF plasmids: cassettes utilizing CaMV sequences for
- 50

expression of foreign genes in plants. J Biotechnol  
14:333-344.

5 van Haaren MJ, Ow DW (1993) Prospects of applying a  
combination of DNA transposition and site-specific  
recombination in plants: a strategy for gene  
identification and cloning. Plant Mol Biol 23:525-533

10 Velten J, Velten L, Hain R, Schell J (1984) Isolation of  
a dual plant promoter fragment from the Ti plasmid of  
*Agrobacterium tumefaciens*. EMBO J 3:2723-2730.

15 Wasmann CC, Reiss B, Bartlett SG, Bohnert HJ (1986) The  
import of the transit peptide and the transport protein  
for protein import into chloroplasts. Mol Gen Genet  
205:446-453

20 Wimmer B, Lottspeich F, van der Klei I, Veenhuis M,  
Gietl C (1997) The glyoxysomal and plastid molecular  
chaperones (70-kDa heat shock protein) of watermelon  
cotyledons are encoded by a single gene. Proc Natl Acad  
Sci USA 94:13624-13629

25 Xiang C, Guerra DJ (1993) The anti-nptII gene. A  
potential negative selective marker for plants. Plant  
Physiol 102:287-293

30 Zoubenko OV, Allison LA, Svab Z, Maliga P (1994)  
Efficient targeting of foreign genes into the tobacco  
plastid genome. Nucleic Acids Res 22:3819-3824

35 While certain of the preferred embodiments of the  
present invention have been described and specifically  
exemplified above, it is not intended that the invention  
be limited to such embodiments. Various modifications  
may be made thereto without departing from the scope and  
spirit of the present invention, as set forth in the  
following claims.

What is claimed is:

1. A site specific recombination method for  
5 removal of predetermined nucleic acid sequences from the  
plastid genome, said method comprising:
  - a) providing a first nucleic acid construct,  
said construct comprising a promoter being operably  
linked to a nucleic acid encoding an optional plastid  
10 targeting transit sequence which is operably linked to a  
nucleic acid encoding a protein having excision  
activity, said construct further comprising a first  
selectable marker encoding nucleic acid having plant  
specific 5' and 3' regulatory nucleic acid sequences;
  - 15 b) providing a second DNA construct, said  
second construct comprising an second selectable marker  
encoding nucleic acid and excision sites, said second  
construct optionally containing a gene of interest, said  
second construct further comprising flanking plastid  
20 targeting nucleic acid sequences which facilitate  
homologous recombination into said plastid genome;
  - c) introducing said second DNA construct into  
a plant cell;
  - d) culturing said plant cell of step c) in the  
25 presence of a selection agent, thereby selecting for  
those plant cells expressing the proteins encoded by  
said second DNA construct;
  - e) introducing said first DNA construct into  
plant cells from step d) in the presence of a selection  
30 agent and selecting those plant cells expressing  
proteins encoded by said first construct, which when  
present said excising activity acts on said excision  
sites, thereby excising said predetermined target  
sequence.

35



2. A method as claimed in claim 1, wherein a plant is regenerated from plant cells of step c), cells are then contacted with said first construct and steps d) and e) are performed.

5

3. A method as claimed in claim 1, wherein said first construct is that depicted in Figure 3.

10

4. A method as claimed in claim 1, wherein said second construct is that depicted in Figure 2.

15

5. A method as claimed in claim 1, wherein said protein having excision activity is selected from the group consisting of CRE, flippase, resolvase, FLP, SSV1-encoded integrase, and transposase.

20

6. A method as claimed in claim 1, wherein said excision sites are LOX sequences, and *frt* sequences.

7. A method as claimed in claim 1, wherein said selection agent is selected from the group consisting of kanamycin, gentamycin, spectinomycin, streptomycin and hygromycin, phosphinotricin, basta, glyphosate and bromoxynil.

25

8. A method as claimed in claim 1, wherein said excision of said predetermined sequence creates an expressible translational fusion protein.

30

10. A method as claimed in claim 1, wherein said predetermined target sequence is the selectable marker encoding nucleic acid present in said second construct.

35

11. A plant regenerated from the method of claim 1.

12. A site specific recombination system comprising the constructs of claim 1.

13. A site specific recombination method for removal of predetermined nucleic acid sequences from the plastid genome, said method comprising:

a) providing a first nucleic acid construct, said construct comprising a regulated promoter being operably linked to a nucleic acid encoding an optional plastid targeting transit sequence which is operably linked to a nucleic acid encoding a protein having excision activity, said construct optionally further comprising a first selectable marker encoding nucleic acid having plant specific 5' and 3' regulatory nucleic acid sequences;

b) providing a second DNA construct, said second construct comprising an second selectable marker encoding nucleic acid and excision sites, said second construct further comprising flanking plastid targeting nucleic acid sequences which facilitate homologous recombination into said plastid genome at a predetermined target sequence such that excision sites flank said predetermined target sequence following homologous recombination;

c) introducing said second DNA construct into a plant cell;

d) culturing a plant cell of step c) in the presence of a selection agent, thereby selecting for those plant cells expressing the proteins encoded by said second DNA construct;

e) regenerating a plant from cells obtained in step d);

f) introducing said first DNA construct into plant cells from step e) in the presence of a selection agent and selecting those plant cells expressing

proteins encoded by said first construct, which when present said excising activity acts on said excision sites, thereby excising said predetermined target sequence.

5

14. A method as claimed in claim 13, wherein said regulatable promoter is selected from the group of promoters consisting of inducible promoters, tissue specific promoters, developmentally regulated promoters and chemically inducible promoters.

10

15. A method as claimed in claim 13, wherein said predetermined target sequence is selected from the group consisting of genes associated with male sterility, clpP ribosomal proteins, ribosomal RNA operon sequences.

15

16. A method as claimed in claim 13, wherein said protein having excision activity is selected from the group consisting of CRE, flippase, resolvase, FLP, SSV1-encoded integrase, and transposase.

20

17. A method as claimed in claim 13, wherein said excision sites are LOX sequences, and *frt* sequences.

25

18. A method as claimed in claim 13, wherein said selection agent is selected from the group consisting of kanamycin, gentamycin, spectinomycin, streptomycin and hygromycin, phosphinotricin, basta, glyphosate and bromoxynil.

30

19. A plant regenerated from the method of claim 13.

20. A site specific recombination system for removal of predetermined nucleic acid sequences comprising the construct of claim 13.

5           21. Progeny plants obtained from the plant of claim 11.

          22. Progeny plants obtained from the plant of claim 18.

10

1/20






Figure 1

2/20



3/20

**Figure 3**




Figure 4





Figure 5

6/20




Figure 6

7/20




Figure 7





Figure 8

9/20



10/20




Figure 10

11/20




Figure 11




Figure 12






Figure 13




Figure 14

15/20




Figure 15

16/20




Figure 16




Figure 17




Figure 18




Figure 19

20/20




Figure 20



## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US00/25930**A. CLASSIFICATION OF SUBJECT MATTER**

IPC(7) : Please See Extra Sheet.

US CL : Please See Extra Sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/69.1, 69.7, 320.1, 410, 419, 468; 800/278, 287, 288, 295, 298, 303

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

WEST, Agricola, Biosis, Caplus

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	LYZNIK et al. FLP-Mediated Recombination Of FRT Sites In The Maize Genome. Nucl. Acids Res. 1996, Vol. 24, No. 19, pages 3784-3789, especially pages 3784-3787.	1-22
Y	DALE et al. Gene Transfer With Subsequent Removal Of The Selection Gene From The Host Genome. Proc. Natl. Acad. Sci. USA. December 1991, Vol. 88, pages 10558-10562, especially pages 10559-10561.	1-6, 10-13, 16, 17, 19-22
Y	BAYLEY et al. Exchange Of Gene Activity In Transgenic Plants Catalyzed By The Cre-Lox Site-Specific Recombination System. Plant Molecular Biology. 1992, Vol. 18, pages 353-361, especially pages 356-358.	1-22



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:	*T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention	
*A	document defining the general state of the art which is not considered to be of particular relevance		
*B	earlier document published on or after the international filing date	*X	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
*L	document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Y	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
*O	document referring to an oral disclosure, use, exhibition or other means		
*P	document published prior to the international filing date but later than the priority date claimed	*A	document member of the same patent family

Date of the actual completion of the international search

06 NOVEMBER 2000

Date of mailing of the international search report

04 DEC 2000

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

ASHWIN MEHTA

Telephone No. (703) 308-0196

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US00/25930

## C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	LYZNIK et al. Heat-Inducible Expression Of FLP Gene In Maize Cells. Plant J. 1995, Vol. 8, No. 2, pages 177-186, especially pages 178-181.	1-22
Y	SHANKLIN et al. The Stroma Of Higher Plant Plastids Contain ClpP and ClpC, Functional Homologs Of Escherichia coli ClpP and ClpA: An Archetypal Two-Component ATP-Dependent Protease. Plant Cell. October 1995, Vol. 7, pages 1713-1722, especially pages 1714-1715 and 1720.	15
Y	SVAB et al. High-Frequency Plastid Transformation In Tobacco By Selection For A Chimeric aadA Gene. Proc. Natl. Acad. Sci. USA. February 1993, Vol. 90, pages 913-917, especially pages 914-915.	1, 13

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US00/25930

**A. CLASSIFICATION OF SUBJECT MATTER:**

IPC (7):

C12N 5/04, 15/09, 15/63, 15/64, 15/65, 15/67, 15/82, 15/87, 15/90; A01H 1/00, 5/00

**A. CLASSIFICATION OF SUBJECT MATTER:**

US CL :

435;/69.1, 69.7, 320.1, 410, 419, 468; 800/278, 287, 288, 295, 298, 303